

DECODING RESISTANT HYPERTENSION SIGNALING PATHWAYS

Ricardo Cambraia Parreira^{a,b}, Leandro Heleno Guimarães Lacerda^c, Rebecca Vasconcellos^{a,b}, Swiany Silveira Lima^a, Anderson Kenedy Santos^a, Vanessa Fontana^d, Valéria Cristina Sandrim^e and Rodrigo Ribeiro Resende^{a,b}

^aDepartment of Biochemistry and Immunology, Institute of Biological Sciences, Universidade Federal de Minas Gerais, Av. Antônio Carlos, 6627, 31270-901 Belo Horizonte, MG, Brazil

^bInstituto Nanocell, Rua Santo Antônio, 420, 35500-041 Divinópolis, MG, Brazil

^cCollege of Sete Lagoas - FACSETE, Rua Itália Pontelo, 50/86, Chácara do Paiva 35700-170 -Sete Lagoas, MG, Brazil

^dDepartment of Molecular & Clinical Pharmacology, Institute of Translational Medicine, Wolfson Centre for Personalised Medicine, University of Liverpool, Liverpool, United Kingdom.

^eDepartment of Pharmacology, Institute of Biosciences of Botucatu, Universidade Estadual Paulista (UNESP), Distrito Rubiao Junior, Botucatu, São Paulo, 18680-000, Brazil.

*Corresponding author: Prof. Dr. Rodrigo R Resende, Cell Signaling and Nanobiotechnology Laboratory, Department of Biochemistry and Immunology, Bloco N4 112, Phone: +55 (31) 3409-2627, e-mail address: rresende@institutonanocell.org.br or rresende@hotmail.com

Abstract Resistant Hypertension (RH) is a clinical condition in which the hypertensive patient has become resistant to drug therapy and is often associated with increased cardiovascular morbidity and mortality. Several signaling pathways have been studied and related to the development and progression of RH: Modulation of sympathetic activity by leptin and aldosterone, primary aldosteronism, arterial stiffness, endothelial dysfunction and variations in the renin-angiotensin-aldosterone system (RAAS). MicroRNAs (miRNA) comprise a family of small noncoding RNAs that participate in the regulation of gene expression at post-transcriptional level. miRNAs are involved in the development of both cardiovascular damage and hypertension. Little is known on the molecular mechanisms that lead to development and progression of this condition. This review aims to cover the potential roles of microRNAs in the mechanisms associated with the development and consequences of RH, and explore the current state of the art of diagnostic and therapeutic tools based on microRNA approaches.

Keywords MicroRNA; Resistant Hypertension; Signaling Pathways

Abbreviations List

20-HETE - 20-hydroxyeicosatetraenoic acid

ABPM - ambulatory blood pressure monitoring

ACE - angiotensin converting enzyme

ADMA - asymmetric dimethylarginine

AGT - angiotensinogen

ALLHAT - antihypertensive and lipid-lowering treatment to prevent heart attack trial

AngI - angiotensin I

AngII - angiotensin II

APCs - angiogenic progenitor cells

AT1 - angiotensin 1 receptor

AVPR1A - arginine-vasopressin receptor

Bcl-2 - B-cell lymphoma 2

BDKRB2 - bradykinin β 2 receptor

bFGF - basic fibroblast growth factor

BP - blood pressure

CAV1 - Caveolin-1

COX-2 - cyclooxygenase-2

CYP4A - cytochrome P450-4A)

EGF - epidermal growth factor

EGFR - Epidermal growth factor receptor

eNOS - endothelial NO synthase

ERK - extracellular regulated kinase

ET-1 - endothelin-1

FOXO1 - forkhead box 1

FoxO1 - forkhead Transcription factor

GATA2 - endothelial transcription factor GATA2

GLUT4- glucose transporter 4

GPCRs - G-protein-coupled receptors

HDL - High density lipoprotein cholesterol

LNA - locked-nucleic-acid

MAPKs - mitogen-activated protein kinases

MCP-1 - monocyte chemoattractant protein-1

MEK - mitogen-activated protein kinase

miRNA – microRNAs

MMP - matrix metalloproteinase

mTOR - mammalian target of rapamycin

NO - nitric oxide

NYHA - New York Heart Association

PAK4 - serine/threonine-protein kinase PAK4

PDGF - platelet-derived growth factor

PG - prostaglandins

PI3K - phosphoinositide 3-kinase

PIK3R2 - phosphoinositide-3-kinase regulatory subunit 2

PIP3 - phosphatidylinositol 3,4,5-trisphosphate

PKC - protein kinase C

PLC - phospholipase C

PLD - phospholipase D

pri-miRNA - primary miRNAs

RAAS - renin-angiotensin-aldosterone system

RH - resistant hypertension

RhoB - ras homolog family member B

RISC - RNA-induced silencing complex

ROS - reactive oxygen species

RTKs - receptor tyrosine kinases

S1P - sphingosine-1-phosphate

Sirt1 - silent information regulator 1

SNS - Sympathetic nervous system

Spred-1 - Sprouty-related protein

SRF - Serum Response Factor

TBXA2R - thromboxane A2 receptor

VCAM1 - vascular cell adhesion molecule 1

VEGF - vascular endothelial growth factor

VSMC - vascular smooth muscle cells

1.Introduction

Resistant Hypertension

Resistant hypertension (RH) is characterized by a condition in which the patient requires four or more antihypertensive medications, including a diuretic, regardless blood pressure control. RH patients can be classified as controlled or uncontrolled according to the achievement of the blood pressure goals (1).

The RH affects approximately 13% to 25% of the hypertensive population (2-4)and represents a risk factor for cardiovascular events. Results from the Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT) showed increased hazard ratios for stroke (1.57), end-stage renal disease (1.95), heart failure (1.88), coronary heart disease (1.44) and all-cause mortality (1.30) in RH compared to non-resistant hypertensive subjects(5).

RH is a multifactorial condition and several environmental and genetic factors contribute to the development and progression of the disease. Firstly, the identification of pseudoresistance must be performed to identify true resistant hypertensive patients, caused mainly due to poor BP measurements, lack of adherence, suboptimal therapy and white coat hypertension. In addition, secondary causes of RH, such as primary aldosteronism, pheochromocytoma, renal artery stenosis or Cushing's syndrome should be identified since the pharmacological treatment will be specific for each condition. Among the factors associated with an increased risk for RH are: older age, African origin, female gender, overweight and obesity (1, 6).

The RH present three relevant characteristics: 1) high incidence of the following comorbidities: obstructive sleep apnea (7), thyroid disorders (7), primary aldosteronism (8, 9), reduced plasmatic renin activity, obesity (6), diabetes *mellitus* (1, 6) 2) high prevalence of target organ damage; and 3) high blood pressure (BP) levels measured by ambulatory blood pressure monitoring (ABPM) (10-12).

The achievement of the BP goals relies on physician examination and on patient characteristics and compliance to pharmacological and non-pharmacological treatments. It is well know that obesity, excessive alcohol and/or salt intake, sedentary lifestyle, smoking, insulin resistance, difficulty in adopting dietary measures and lack of adherence to therapeutic treatment affects BP control (12). In addition, prescription of

high-cost medicines by physicians, multiple administration regimens, suboptimal doses and presence of adverse effects are associated with uncontrolled BP (13).

The factors associated with diagnostic and treatment of RH, including lifestyle, detailed history of medication adherence, correct BP measurement, biochemical analysis for dosage of electrolytes, glucose and creatinine, as well as determination of protein and sodium in the urine (12).

It is also necessary the exclusion of pseudohypertension. For example, Mönckeberg's sclerosis is a condition characterized by the loss of elasticity and thickening of the walls of the muscular arteries caused by a calcification of the tunica media constituted of smooth muscle. In the measurement of blood pressure by non-invasive methods, the patient presents high BP values, while in reality its pressure is normal, therefore, it is necessary to use invasive measurement methods to correctly measure its BP (12). White coat hypertension, also a condition of pseudoresistance, where the patient exhibits high values during the verification in the physician's office. It can be excluded by 24 hours ABPM. It is estimated that 30% of patients with elevated BP and with treatment of up to 3 drugs present this condition of white coat hypertension (12, 14).

In order to complete the diagnosis of RH, there are some clinical situations that are considered as secondary causes of this condition, such as: primary hyperaldosteronism, pheochromocytoma, fibromuscular dysplasia, patients with increased risk of atherogenesis (12), obstructive sleep apnea, renal artery stenosis, renal parenchymal disease, Cushing's syndrome, and thyroid and parathyroid diseases (10). These forms of secondary hypertension present high prevalence in association with RH, as can be verified in table 1. Hyperaldosteronism results from the excessive production of aldosterone, a hormone that is produced in the adrenal glands and that decreases the excretion of sodium and increases the excretion of potassium by the kidneys, sweating, and saliva. The determination of the rate of aldosterone/renin ratio is used as screening tool for hyperaldosteronism diagnostic, and if any alteration is confirmed, diagnostic imaging and blood samples from each side of the adrenal glands are used to corroborate the diagnosis (12). Obstructive sleep apnea consists of the collapsing of the pharynx walls hampering the adequate respiration of the individual. The patient is then submitted to a nocturnal polysomnography in order to monitor respiration and body functions

duringsleep for diagnostic of the condition (15). For more information on diagnostic and treatment methods, see table 1 (14) .

A flowchart (**Figure 1**) summarizes the steps involved in the diagnostic of RH according to the American Heart Association Statement (1, 16).

[TABLE 1]

[FIGURE 1]

Although RH is a multifactorial condition, sodium excess, fluid retention, increased activation of the Renin Angiotensin Aldosterone System and higher sympathetic tone are among the most well-described mechanisms of BP elevation in RH. The complex pathophysiology of the development and progression of RH requires further investigation to identify molecular mechanisms that could be translated into diagnostic and more assertive therapeutic strategies. In the following sessions, the signaling pathways and the participation of miRNA in their regulation will be discussed.

2. Renin Angiotensin Aldosterone System

The Renin-Angiotensin-Aldosterone System (RAAS) is responsible for the hemodynamic equilibrium. This is possible due to the effects of this system on the kidneys, that act in the sodium water balance, and also due to the influence in the vascular resistance on the peripheral blood vessels, thus permitting the maintenance of the BP (17). In order to the RAAS to produce a response it is necessary that occur some type of alteration in the circulating blood volume, such as blood loss, dehydration or even pumping failure by the ventricles (18) (**Figure 2**).

Juxtaglomerular apparatus comprises afferent arterioles in the distal part of the ascending branch of the loop of Henle in the renal glomeruli. The cells that line these arterioles in the region of the apparatus are called juxtaglomerular cells and are able to recognize the BP inside these vessels (18). Moreover, the cells that line the loop of Henle in the region of the juxtaglomerular apparatus are called macula densa and respond to changes in the sodium concentration of the filtrate. By detecting these changes, the cells of the dense macula stimulate juxtaglomerular cells to produce an enzyme called renin that is released into the bloodstream. This enzyme is responsible for the production of angiotensin I (AngI) through the cleavage of angiotensinogen, which is synthesized and secreted by the liver. In the pulmonary and renal endothelium, an enzyme called angiotensin converting enzyme (ACE), which hydrolyzes the circulating AngI in angiotensin II (AngII), is present. The angiotensin 1 receptor (AT1) is activated by AngII thus promoting a vasoconstrictive response in the blood vessels, in addition to stimulating the adrenal gland to produce aldosterone. The renal tubules respond to the aldosterone by retention of sodium and water, which will promote increased blood volume and consequently, BP elevation (19, 20).

[FIGURE 2]

The RAAS constitutes the main signaling pathway involved in the long-term control of the BP. Innumerable regulators participate in this biochemical cascade of communication such as renin, angiotensinogen, AngI and II, ACE 1 and 2, aldosterone and angiotensin- (1-7) [Ang- (1-7)]. The deregulation of one or more effectors of this system contributes to failures in blood pressure control, usually leading to hypertension (21). Resistant Hypertension is accompanied by intravascular fluid retention that can be attributed, at least in part, to dysregulation in the renin-angiotensin-aldosterone system. Previous studies have found evidence of intravascular volume expansion (higher levels of brain-type natriuretic peptide – BNP, and atrial natriuretic peptide - ANP) and aldosterone excess (higher levels of plasma and urinary aldosterone, aldosterone to renin ratio) in resistant hypertension compared to controls (22). Similarly, another study reported higher volume of fluid by thoracic electrical bioimpedance, suggesting that the intensification of diuretic therapy in those patients could be beneficial (23). ANP and BNP are hormones that regulate cardiovascular hemodynamics. They are secreted by, respectively, cardiac atria and cardiac ventricles, in response to stretch or pressure. Natriuretic effects are mediated by subtype A-natriuretic peptide receptor, which is expressed in several tissues, including kidneys, blood vessels, adrenal glands, and adipose tissue. ANP produces its natriuretic actions by increasing glomerular filtration rate, inhibits sodium transport in proximal tubule and inhibition of aldosterone release in adrenal cells. The latter effect is also attributed to BNP (24). Aldosterone is one of the most studied RAAS components in RH, several studies had shown that aldosterone excess is a common characteristic of RH. In addition, primary aldosteronism is the most common secondary cause in the patients with RH (8). This condition is characterized by excessive autonomic secretion of aldosterone by the adrenal gland, being the production of adenomas and idiopathic hyperaldosteronism the main forms (32, 33). This secretion, stimulated by renin, promotes the retention of sodium and water, promoting the elevation of blood volume, and consequently the increase of BP. When it is released into the bloodstream, the aldosterone diffuses through the membrane into the cytosol of renal tubular epithelial cells, subsequently binding to a family of NRC2-type mineralocorticoid receptors. This aldosterone-receptor complex will be translocated to

the nucleus, activating the synthesis of proteins related to sodium and potassium transport, such as Na⁺ K⁺ ATPase (**Figure 3**) (25).

[**FIGURE 3**]

An important signaling pathway in the primary aldosteronism is the PI3K (phosphoinositide 3-kinase) pathway with the activation of mTOR (mammalian target of rapamycin), when overactivated is involved in the tumorigenesis and metastasis in some types of human tumors such as renal cancer, adrenal carcinoma and pheochromocytoma (26-28) (**Figure 4**) (27). The PI3K/AKT/mTOR pathway is regulated in response to the signaling of growth factors such as the epidermal growth factor (EGF) through receptor tyrosine kinases (RTKs)(29).

Another stimulus to the release of aldosterone is the action of sphingosine-1-phosphate (S1P) by means of the activation of the PI3K/AKT (protein kinase B) and MEK (mitogen-activated protein kinase)/ERK (extracellular regulated kinase) pathway in glomerular cells of the adrenal glands (**Figure 4**) (28). S1P is a bioactive sphingolipid intracellularly formed that acts as a second messenger mediating regulatory processes such as cell differentiation, migration and contraction, modulation of immune response and angiogenesis, and this molecule is considered to be the key hormone for hemodynamic stability in humans (30, 31). Its action involves the activation of phospholipase D (PLD), calcium influx (Ca²⁺) from the extracellular medium and phosphorylation of α and β isoforms of protein kinase C (PKC) (30, 32, 33).

[**FIGURE 4**]

Previous studies had shown excess of aldosterone in uncontrolled RH compared to controlled group. The same study demonstrated that aldosterone was correlated to arterial stiffness (34). Furthermore, higher aldosterone levels were associated with the T allele for the polymorphism -344 C/T *CYP11B2* (aldosterone synthase gene) in RH

subjects, and this effect was shown to be more pronounced in patients under spironolactone treatment (35). Studies have demonstrated significant reductions in blood pressure with addition of mineralocorticoid receptors antagonists, such as spironolactone, and that drug has been suggested as the optimal fourth-line drug for BP control in RH (159).

Angio-(1-7) is a heptapeptide that carries out an important function in the RAAS. This molecule is formed both by the action of ACE 1 (dependent pathway) and the hydrolysis of AngII by the ACE 2 (independent pathway) (36, 37) being this last one the most important pathway in the formation of Ang-(1-7) (38). This molecule produces its AngII endogenous counter-regulatory effects on RAAS (vasodilation, cardio protection, natriuresis and diuresis, angiogenesis inhibition, and cellular growth) (39) through the binding to its specific receptor called Mas, a G protein-coupled receptor (32, 40, 41).

The ACE2/Ang-(1-7)/Mas signaling pathway consists in one of the RAAS axes that opposes, in terms of function, to another classical axis of this system, the ACE/AngII/AT₁R. The imbalance of these two opposing axes, mainly in the direction of the ACE/AngII/AT₁R axis predisposes to cardiovascular diseases and other disorders (37, 41).

The Ang-(1-7)/Mas complex regulates different signaling pathways, such as PI3K/AKT and ERK signal, and involves the maintenance of some effectors like nitric oxide (NO)(32, 41), FOXO1 (forkhead box 1)(39) and cyclooxygenase-2 (COX-2)(40) (**Figure 5**). Studies report that due to the participation of the Ang-(1-7) in these mechanisms, this heptapeptide is related to pathological conditions such as fibrosis and inflammatory processes in some organs, like lungs, liver and kidneys (42). Other findings demonstrate that Ang-(1-7), through the interaction with its Mas receptor, stimulates the activation of the nitric oxide synthesis (eNOS) in endothelial cells, promoting vasodilation (32, 36, 41).

Another study demonstrates that Ang-(1-7), through the interaction with its specific Mas receptor promotes the increase of nitric oxide (NO) and prostaglandins (PG) synthesis and release leading to vasodilation and inhibition of cellular growth, opposing to the vasoconstrictor and proliferative effects mediated by the interaction of AngII with its AT₁ receptors. The imbalance between these two axes of the RAAS,

reflected by the imbalance between these peptides, which are observed in cardiovascular diseases, can lead to the decrease of NO and consequently to endothelial dysfunction (**Figure 5**) (43).

[FIGURE 5]

2.3. Sympathetic Nervous System

Sympathetic nervous system regulates cardiac output and peripheral vascular resistance (vasoconstriction) through release of norepinephrine and epinephrine, resulting in increase in blood pressure. At the renal level, SNS activation increases renin release from juxtaglomerular cells and modulate tubular sodium reabsorption (44). RH patients have reduced heart rate variability, which is a marker for SNS activity. It was shown that 63% of the patients present a nondipping pattern (BP do not drop at night), which indicate sympathetic overflow. Moreover, sympathetic activation also increase sodium reabsorption and promote renin secretion and renal denervation has been investigated for treatment of resistant hypertension. In spite first studies in humans had shown promising results, randomized and blinded clinical trials demonstrated no benefit on BP control compared to sham procedure (45). Another intervention that has been tested is baroreflex activation therapy. Carotid sinus stimulation reduces BP in patients with uncontrolled RH, showing the import role of sympathetic activity in this condition (46-48).

2.4 - Adipokines

Adiponectin and leptin are two of the adipokines produced in adipose tissue. Obesity is an important comorbidity in RH and plasma levels of adiponectin and leptin were reported to be lower and higher, respectively in RH (34) (49). Leptin is a peptide hormone that is expressed in a variety of tissues, such as lymphoid tissue, pituitary gland, skeletal muscle, placenta, and ovary (50). However, white adipose tissue is the main responsible for the synthesis and secretion of this peptide, which has as effects to act on the hypothalamus in order to decrease appetite and stimulate sympathetic activity of the nervous system (**Figure 6**) (51). Elevated levels of leptin stimulate neurons in the hypothalamus to secrete a precursor protein that is cleaved in α -melanocyte stimulating hormone, which binds to melanocortin 3 and melanocortin 4 receptors. The binding of this peptide to the receptors stimulates the sympathetic nervous system, elevates the energy expenditure, decreases food intake, and activates the hypothalamic-pituitary-adrenal axis (52-54). A mechanism that demonstrates which factors are involved in the generation of hypertension associated with obesity is represented below (**Figure 6**) (51).

[**FIGURE 6**]

2.5. Insulin resistance and hypertension: the role of the Caveolin-1 (CAVI) gene

It was recently demonstrated that the gene Caveolin-1 (*CAVI*), located in the chromosome 7q31.1 (55) constitutes a gene that is associated with metabolic dysfunction in animal and cellular models, especially in insulin resistance, proving to be a potential marker for this condition in human beings (56). Genetic variations in *CAVI* are involved in the mechanism of insulin signaling and vascular function (**Figure 7**), shown in studies with animal models and cell culture (57, 58).

Increase *in* homeostasis model assessment *of* insulin resistance (*HOMA-IR*), showing that *CAVI* is not only a genetic marker for dysfunction but also provides information about a potential mechanism of development of insulin resistance and hypertension in humans (56).

CAVI is a regulatory gene for insulin signaling and insulin receptor stability (56). Specifically, *CAVI* binds directly to the insulin receptor in the adipocytes and the disturbance of this complex by GM3 ganglioside causes alteration in insulin signaling

(59) (**Figures 7 and 8**). In addition, the decrease in *CAVI* activity results in a 90% reduction in insulin receptor levels in the adipocytes of knockout rats (60).

Although the role of *CAVI* in insulin-mediated glucose uptake is not well elucidated (60), this gene demonstrated relevance in the translocation of glucose transporter 4 (GLUT4) to adipocyte (61) and muscle cells (62).

[**FIGURE 7**]

[**FIGURE 8**]

2.6 Vascular stiffness and endothelial dysfunction

Previous studies has been showed the participation of vascular stiffness and endothelial dysfunction in the pathogenesis of resistant hypertension. An increased carotid-femoral pulse wave velocity was observed in RH patients compared to non RH hypertension, demonstrating the impairment of elasticity in these vessels. In addition, flow-mediated dilation was found to be reduced in RH, reflecting an endothelial dysfunction (63).

2.6.1 Epidermal growth factor receptor in the vascular smooth muscle

The activation of the signaling pathway of the EGFR by matrix metalloproteinase (MMP), stimulated by GPCR agonists, such as catecholamines, endothelin-1 (ET-1) and AngII lead to the increase of the oxidative stress, promote stimulation of the hypertrophic growth and consequently increase of the muscular tone in hypertension (**Figure 9**) (64). Among these receptors it can be mentioned adrenoceptors and angiotensin receptors that contribute to the hypertension pathogenesis mainly through the vasoconstrictor effects produced after stimulation (65, 66).

The vasoconstrictor responses promoted by this pathway are mediated by phospholipase C (PLC), DAG and Ca^{2+} besides the growth promotion pathway

involving the tyrosine receptor and mitogen-activated protein kinases (MAPKs) (**Figure 9**) (67). Studies have shown a connection between the GPCR stimulus with the MAPK signaling pathway (through the dependent activation of MPM) in the vascular smooth muscle cells (65, 68-70). Associations have been made between GPCR stimulus by MPM's such as MPM-2, MPM-3, and MPM-7 in cardiomyocytes, fibroblasts and epithelial and endothelial cells (71) with consequent development of cardiovascular hypertrophy associated with hypertension (72-75).

[**FIGURE 9**]

2.6.2 CYP4A (cytochrome P450-4A)/20-HETE (20-hydroxyecosatetraenoic acid)

20-HETE is an arachidonic acid metabolite formed through reactions catalyzed by the cytochrome P450-4A enzymatic complex (CYP4A) in vascular smooth muscle cells and is related to vascular dysfunction and arterial hypertension (**Figure 10**) (76). This molecule has vasoconstrictive action and exerts an important role in vascular function and in the development and progression of cardiovascular diseases (77). Studies have demonstrated the relationship between genetic variations in precursors of 20-HETE formation and the elevation of this metabolite and the BP in humans (78, 79).

In Dahl SS (salt-sensitive) rats, a genetic model of salt-sensitive hypertension, 20-HETE has been shown to contribute to the increase of total peripheral resistance by reducing the ability of the vascular system to respond to direct vasodilation stimulation by reducing vascular function, thus contributing to an increase in BP (80, 81).

Some studies demonstrate that ROS (reactive oxygen species) are important molecules in the development of oxidative stress, playing an important role in vascular dysfunction in Dahl SS rats (82, 83). The chronic exposure to low levels of AngII in these animals may lead to an increase in oxidative stress by elevating ROS cellular concentrations, thus contributing to the reduction of vascular relaxation even when these animals are submitted to a sodium restriction diet or are normotensive (84).

[**FIGURE 10**]

3. MicroRNAs

MicroRNAs (miRNA) are members of a class of small non-coding RNAs that are able to interfere with the translation of several genes (85). They are highly conserved molecules present in many species and components of the miRNA synthesis machinery can be found in bacteria and species of archaea scattered (86). The miRNAs are single stranded molecules formed of approximately 22 nucleotides, generated from endogenous transcripts (86), expressed from intergenic regions or from introns. The biogenesis of miRNA begins in the nucleus from long precursors called primary miRNAs (pri-miRNA) transcribed by polymerase II (87). The pri-miRNAs, still in the nucleus, are processed by *Drosha* and its cofactor DGCR8 (88, 89), that remove a region of the double-stranded ribbon and flanking sequences resulting the pre-miRNAs. The pre-miRNAs are transported to the cytoplasm via Exportin-5 (90). In the cytoplasm, the pre-miRNAs are cleaved by a ribonuclease III, called *Dicer*, forming the mature miRNAs (91). The mature miRNA associates to a complex of enzymes called RISC and acts on the translation of mRNA according to the complementarity between the miRNA and its target (92, 93) (**Figure 11**). A complete pairing between the miRNA and its target causes cleavage of the mRNA while an incomplete pairing causes the silencing of the target mRNA expression (94).

The miRNAs are among the most abundant gene regulatory molecules (95) and participate in important regulatory cell pathways, such as apoptosis, cell proliferation, differentiation and development (96-98), and also in various cardiovascular conditions such as cardiac hypertrophy, arrhythmias, apoptosis, and regeneration of cardiomyocytes, fibrosis, heart failure (99), coronary heart disease, acute myocardial infarction (100) and essential hypertension (101).

Besides, miRNAs are currently considered as new hormones or molecules that carry information through intercellular mechanisms. they can be transported by body fluids, such as blood, serum, urine, breast milk, circulating extracellular vesicles, exosomes and microvesicles, in addition to complexes with particles of HDL, protected from degradation by RNases and other enzymes (102), thus being characterized as potential molecular biomarkers for disease prognosis or as a predictor of therapeutic efficacy (103).

[FIGURE11]

3. 1. miRNAs involved in hypertension

Due to the difficulty of studying RH in animal and *in vitro* models, the studies are clinical and poorly understood in terms of mechanisms. Thus, we decided to describe the microRNAs that interact with the most relevant pathways in RH. To date, several miRNAs have been identified and are related to the complications of resistant hypertension (**Figure 12**). The most important are listed below (Table 2):

[TABLE 2]

1. RAAS

The miRNAs miR-181a and miR-663 appear to be directly related to the hypertensive condition. Patients with untreated hypertension demonstrated a low expression of miR-181a and miR-663 in kidney samples. Experiments in culture demonstrated that miR-181 and miR-663 were able to regulate renin expression. The data show that the low expression of these two miRNAs causes the elevation of the renin expression leading to the hypertensive condition (104). In addition, the presence of miR-181a in the serum of hypertensive patients was verified. Little is known about the role of miR-181a in the control of renin or whether there are other targets in the kidney. Future work should focus on the regulatory role of miR-181 over the other components of RAAS (105).

The expression of AT₁R (angiotensin II type I receptor) and AT₂R (angiotensin II type 2 receptor) are regulated mainly by post-transcriptional mechanisms. It was verified that miR-155 (expressed in fibroblasts and VSMCs) and miR-802 (expressed in intestinal epithelial cells), in functional assay, are responsible for the repression of AT₁R expression, not by decreasing levels of mRNA, but by blocking mRNA translation. Consequently, it leads to the reduction of activation of Angiotensin II-induced ERK1/ERK2 signaling pathway (106-108). Li et al. demonstrated the existence of a pattern of expression of miRNAs in the plasma of hypertensive patients when compared with controls. In this study, an increase in the expression of miR-296-5p, let-7e, hcmv-miR-UL1 and hcmv-miR-UL112 was verified, which acts via IRF-1 (109). There are evidences that the IRF-1 transcription factor up-regulates the AT₂R and plays a role in angiogenesis, neointima formation, endothelial function and SMC migration (109).

An imbalance in the angiotensin II signaling pathway may lead to hypertensive condition. miR-483-3p is a VSMC-specific miRNA and demonstrates to be a general regulator of RAAS, since several components of this system contain the target sequence for this miRNA. Experiments in cell culture proved that this miRNA acts by suppressing the protein levels of angiotensinogen (AGT), angiotensin I converting enzyme (ACE-1), angiotensin II converting enzyme (ACE-2) and AT₂R without altering the levels of its transcripts (110).

Reports show that miR-132 and miR212 are involved in angiotensin II-induced hypertension. Hypertensive rats show elevated expression of miR-132 and miR-212 in the heart, aortic wall, kidneys and plasma after 10 days of sustained hypertension by angiotensin II infusion. The treatment of hypertensive patients with AT₁R receptor blockers exhibits a lower expression of miR-132 and miR-212 confirming their influence on the hypertension phenotype. miR-132 and miR-212 act by activating the Gαq/ ERK1/ 2 signaling pathway (111). In a case-control study, 9 single nucleotide polymorphisms (SNPs) were genotyped in 7 RAAS-related genes with a common site of miRNA recognition in the 3'-UTR region of the mRNA of these genes. In order to verify the functionality of each SNP in the mRNA/miRNA interaction a dual-luciferase reporter gene was used. Four SNPs located in genes associated with blood pressure were reported: the gene coding the arginine-vasopressin receptor (AVPR1A), the gene coding the bradykinin β2 receptor (BDKRB2), and the gene coding the thromboxane A2 receptor (TBXA2R) (112).

Nossent et al. demonstrated that rs11174811 SNP in AVPR1A interferes in the interaction of this gene with miR-526b and miR -578 and, as a consequence, there is an increase in the expression of vasopressin receptor (112) causing an increase in blood pressure. rs5225 and rs2069591 SNPs in BDKRB2 interfere in the interaction of these genes with miR-34a and miR-34c, culminating in an increase of this receptor expression, consequently inducing vasodilation and inhibiting re-uptake by aquaporin. miR-151-3p inhibits the expression of the bradykinin receptor interacting with the rs2069591 SNP located in the PTK2B gene which, in turn, has its vasopressin expression stimulated. miR-765 interacts with rs13306046 SNP of thromboxane A2 receptor, a potent vasoconstrictor, causing an inhibition of the expression of this receptor and a consequent decrease in blood pressure. miR-383 has been associated with increased risk of myocardial infarction but is not associated with the increase in blood

pressure. This miRNA acts on the rs5534 SNP repressing the NR3C2 gene coding the mineralocorticoid receptor (aldosterone receptor) (112).

In the same way, the polymorphisms encountered in the 3'UTR region of SLC7A1 mRNA (solute carrier family 7 member 1), which has more or less sites for miR-122 binding causing a differential expression of it, leading to hypertension phenotype with endothelial dysfunction with decrease of nitric oxide and L-arginine metabolism phenotype (113-115).

The miRNA 24 is expressed in the adrenal cortex and regulates the production of aldosterone and cortisol, since it binds to the 3'UTR region of the mRNAs CYP11B1 (11 β -hydroxylase) and CYP11B2 (aldosterone synthase) which repress them, leading to the decrease in the production of these hormones. Deregulation of production may lead to hypertension. This same miRNA is expressed in cells in the lung where hypoxia leads to the increase of miR-24 expression and the activation of smooth muscle cells (116). Intracellular raise of miR-24 leads to diminished expression of the GATA2 (Endothelial transcription factor GATA2) and PAK4 (Serine/threonine-protein kinase PAK4) transcription factors with functions defined in vascular biology (117) and its inhibition induces the increase of angiogenesis (118).

In experiments with luciferase it was proved that miR-124 and miR-135a are associated with hypertension and they interfere on expression of the nuclear receptor of subfamily 3, group C, member 2 (NR3C2), an aldosterone receptor that participates in the control of blood pressure by promoting the retention of salt by the kidney. miR-124 and miR-135a reduce the protein expression of NR3C2, contributing to the modulation of aldosterone signaling in RAAS, reflecting in the control of blood pressure (119).

miR-584 and miR-31 are predicted to act in a 3'UTR region on 11525C SNP in the human angiotensinogen (AGT) gene. In a cell culture study, the transfection of these miRNAs reduced the levels of AGT expression. Patients that presented 11525A SNP of AGT gene have higher plasma and tissue AGT levels, and consequently increased blood pressure due to the lower interaction of this gene with miR-584 and miR-31 when compared to patients with 11525C SNP (120).

Studies show that miR-145 and miR-143 are essential for differentiation (121) and modulation of vascular smooth muscle cells (VSMC), since the elimination of Dicer

in these cells causes profound hypotension, reflecting *in vivo* contractile dysfunction, blood pressure regulation defects, and vascular remodeling (122). miR-143 and miR-145 demonstrated a relationship to the fate of the VSMCs and maintenance of contractility since KO rats for these two miRNAs exhibit a phenotype that is similar to the rats in which Dicer was eliminated. In addition to this, the overexpression of miR-145 in KO animals for Dicer regains vessel contractility (122).

Boettger et al. demonstrated that the expression of miR-143 and miR-145 during mouse development is essential for the contractile phenotype of VSMC (123). Xin et al. showed that the Serum Response Factor (SRF) controls the expression of miR-143 and miR-145 in VSMC and the absence of these miRNAs causes an imbalance in the feedback cycles needed for the cytoskeleton homeostasis in the VSMC compromising the contractile phenotype of these cells (124). miR-145 acts by suppressing ACE and, in this way, by controlling the stimulation of angiotensin II in VSMCs, thus contributing to the preservation of its contractile phenotype and maintenance of normal blood pressure (123).

miR-143 and miR-145 knockout rats, the actin stress fibers are in disorder, compromising the migratory activity of VSMCs (124), which impair the formation of neointima in response to vascular injury and overregulates (121) the ACE in VSMC (123). There are evidences that these miRNAs are closely involved in the phenotypic reversion of VSMCs during hypertension (121).

In hypertensive patients, miR-9 levels are significantly inferior compared to normotensive patients (125). miR-9 is reported to be a negative regulator of cardiac hypertrophy (126) through inhibition of myocardin. Myocardin is a protein that is found in elevated expression in patients with cardiac hypertrophy and activated by the binding of nuclear factor of activated T-cells c3 (NFATc3) to its promoter region (126).

mir-126 is highly expressed in hypertensive patients (138), which favors angiogenesis and vascular integrity *in vivo* (127, 128). Studies have shown a correlates between miR-126 and Sprouty-related protein (Sprd-1) repression (127, 128). Sprd-1 interacts and inactivates RAF1 negatively regulating the signaling of the Vascular endothelial growth factor (VEGF) through MAP kinase, by repression of the phosphoinositide-3-kinase regulatory subunit 2 (PIK3R2), a negative regulator of PI3K, another pathway activated by the signaling of the VEGF and with vascular cell adhesion

molecule 1 (VCAM1) repression, limiting the adherence of leucocytes to the endothelial cells (129).

miR-21 is significantly upregulated in hypertension (130) and unlike miR-126, miR-21 acts as a negative modulator of angiogenesis inhibiting the proliferation, migration and the ability to form tubes from endothelial cells and is down-regulated by serum and bFGF. It was observed that the overexpression of miR-21 decreases Ras Homolog Family Member B (RhoB) expression and activity, which appears to be associated with migration and tubulogenesis of endothelial cells (131).

miR-21 is activated by asymmetric dimethylarginine (ADMA), which interferes with angiogenesis by direct inhibition of eNOS, causes dysfunction in the circulation of angiogenic progenitor cells (APCs), represses the expression of superoxide dismutase 2, a key enzyme in the defense of oxidative stress, and inhibit SPRY2, causing MAP kinase ERK activation, resulting in the decrease of NO availability and APC dysfunction (132). After angioplasty, in the vascular wall, there is a significant increase in miR-21 that partially reduces the expression of Phosphatase and tensin homolog (PTEN), a phosphatidylinositol 3,4,5-trisphosphate (PIP3) enzyme that inhibits the PI3K pathway by dephosphorylating PIP3 and consequently prevents the activation of AKT (133) and increases the expression of B-cell lymphoma 2 (Bcl-2) (134). Inhibition of miR-21 has a negative effect on neointima formation on vascular injury after angioplasty, decreasing cell proliferation and increasing apoptosis (134). They favor the proliferation of SMCs in culture, inhibiting p27 (Kip1) and p57 (Kip2), which are cell cycle inhibitors, and are upregulated in rat carotid arteries after angioplasty and in smooth muscle cells after vascular wall injury (135, 136). Besides that, it was verified that miR-222 acts by suppressing the expression of miR-221 (135). The treatment of VSMC with platelet-derived growth factor (PDGF) increased miR-221 transcription proving to be critical for the de-differentiation of VSMC, a characteristic of intimal hyperplasia, and caused a decrease in c-kit, a critical gene for the expression of Myocardin (Myocd) and activation of genes in VSMC, modulating the contractile phenotype of these cells. In this study, miR-221 acts via p27 (Kip1) (136).

During aging, there is a progressive increase in the expression of miR-217 in endothelial cells in the same time there is a decrease in the expression of SirT1 (silent information regulator 1), a NAD-dependent deacetylase that controls transcription factors such as FoxO1 (forkhead Transcription factor), p53 and PGC-1 α (peroxisome

proliferators activated receptor gamma coactivator-1a). SirT1 is involved in angiogenesis, longevity and resistance to diseases. It was verified that this miRNA represses the expression of SirT1 blocking the angiogenesis. Another miRNA, miR-34a is also related to the control of senescence (137).

miR-29b is the main regulator of several genes related to collagen synthesis besides others related to the production of extracellular matrix which are determinants in renal injury, its suppression causes the increase of the expression of different types of collagens in the renal cortex causing fibrosis of the renal marrow (138). In the heart, the inhibition of miR-29b is associated with cardiac fibrosis and its increased expression inhibits this phenotype. The decreased in the expression of this miRNA is induced by angiotensin II via TGF- β /Smad3 (139).

As exemplified above, miRNAs play a central role in the fine regulation of the RAAS pathway and when expressed in abnormal conditions can cause a number of cardiovascular effects associated with hypertension.

2. Sympathetic

Renal sympathetic activity appears to be essential in maintaining hypertension. Genetically hypertensive mice (BPH/2J) show hyperactivity of the SNS and exacerbate differences in BP during the day and night periods are associated with a greater neuronal activity. Jackson et al. demonstrated that acute sympathetic ganglion blockade with ACE inhibition by enalapril resulted in greater BP declines in BPH/2J animals when compared to normotensive animals, and thus BPH/2J mice presented a sympathetic contribution of up to 4 fold in the elevation of BP. In these animals, renal renin mRNA levels are upregulated and appear to be regulated by miR-181a, which is found to be lower in BPH/2J animals when compared to normotensive animals (140). After bilateral renal denervation, BPH animals show a greater drop in BP after administration of enalapril, lower levels of miR-181a and greater renin expression compared to normotensive animals (141).

Dorr et al. evaluated the effects of renal sympathetic denervation on serum levels of miR-133a. They found elevated levels of miR-133a in the plasma of hypertensive patients 6 months after renal sympathetic denervation demonstrating a correlation between systolic blood pressure and the renal sympathetic nervous system (142).

3. Adipokines

To date, few miRNAs have been associated with hypertension regulation through the adipokine pathway in humans, also correlated with obesity and diabetes. Recently, miR-199a-3p expression has been found to increase during adipocyte cell differentiation and is upregulated in obese individuals. The regulation is mediated by adipokines (leptin, TNF- α , and IL- β increase expression, while resistin decreases expression) in mature adipocytes. This miRNA possesses as possible NLK (nemo-like kinase), VAMP3 (vesicle-associated membrane protein 3) and Cdk7 (cyclin-dependent kinase 7) targets. Thus, it is speculated that this miRNA is related to the modulation in phenotypes of insulin resistance, inflammatory response and diabetes in obese individuals (143).

In a case study with diabetic patients new potential modulators for the adipokine pathway were discovered. Let-7a and let-7f microRNAs were downregulated and miR-326 was upregulated. The latter possesses adiponectin, adiponectin receptor (ADIPOR)-1, ADIPOR-2, and APPL-1 (molecule involved in the intracellular pathway of insulin signaling) as possible targets. In addition, there was an inverse correlation between the concentration of this miRNA and adiponectin. When treated for glycemic control, patients present higher serum levels of let-7a and let-7f microRNAs, but no significant change was observed for miR-326 (144).

In type 2 Diabetes Mellitus, miR-126 is also decreased (145). Zampetaki et al. demonstrated that, in Diabetes Mellitus, elevated glucose levels cause the reduction of miR-126 in apoptotic bodies of endothelial cells, which can cause less delivery of this miRNA, through these apoptotic bodies or microparticles, to the monocytes and in this way increase the resistance to VEGF and contribute to endothelial dysfunction (145).

4. Therapies involving miRNAs

The miRNAs are excellent candidates for therapeutic agents, since that in many diseases, including in hypertension, miRNA levels are altered. This therapeutic tool can be used to re-establish levels of a miRNA and therefore the lost function as demonstrated by Zhang et al. (2014) in which the administration of miR-29b in animal

model of angiotensin II-induced hypertension blocks progressive heart injury demonstrating its function in chronic coronary disease protection (139).

In recent years, the most used strategy is to inhibit inappropriate expression of an altered miRNA (146). For this to happen, chemically modified oligonucleotides, called antagomirs or anti-miRs, were developed which are administered intravenously and are capable of silencing specific endogenous microRNAs (101).

Nevertheless, we still encounter problems regarding the delivery of these oligonucleotides, since a single miRNAs can interact with several RNAs and, depending on the mode of delivery, on different cell types (147). For example, systemic administration of anti-miR and anti-miR-21 blocked the development of cardiac fibrosis but it was also reported to have effects on the development of renal and pulmonary fibrosis (101). To avoid this situation and increase the efficacy of anti-miRs delivery, various delivery modes of such anti-miRs are being tested, though viral carriers such as plasmids, adenovirus, adeno-associated virus, retrovirus and lentivirus, or nonviral vehicles, of which we can highlight the liposomal delivery system, which is a lipid-based nanoparticle (148), such as cholesterol or conjugated to gold nanoparticles, for example (101, 146).

Recently novel classes of miRNA inhibitors are being produced, such as locked-nucleic-acid (LNA), which exhibits greater stability and prolonged storage inside cells by adding a methylene bridge between the 2'-O and the 4'-C atoms in the ribose ring (123); and the artificial sponges of miRNAs, which are produced from transgenes within cells, and bind to miRNA production sites, blocking the transcription of the entire miRNA family (101).

The miRNAs have great potential as biomarkers of innumerable diseases. The miRNAs express a reproducible pattern in biological fluids in hypertensive patients (95, 123) and are easily isolated from plasma, urine, saliva and amniotic fluid (95) without suffering enzymatic degradations and changes due to changes in temperature and pH (102).

[FIGURE 12]

CONCLUDING REMARKS AND FUTURE DIRECTIONS

This review aimed to compile the knowledge generated by basic and applied sciences mainly regarding microRNA studies. These may help health professionals improve diagnostic and therapeutic accuracy, reducing mortality and morbidity in RH.

Epigenetic as well as genetic factors are identified every day and they are associated with variation in blood pressure levels. As reviewed herein, mutation-polymorphism in some signaling pathway gene may increase or decrease the expression of some microRNAs which are involved both in RH development, therapy response as RH associated complications such as renal failure, coronary artery disease, cardiac hypertrophy, stroke among others. Therefore, the use of microRNA as biomarkers in prevention, diagnosis and therapy of this disease, may help to understand the disease, improve pharmacology therapy as well prevent complications.

Author Contribution

All authors were involved in writing the paper and had final approval of the submitted and published versions.

Funding

This work was supported by Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP), Fundação de Amparo à Pesquisa de Minas Gerais (FAPEMIG), Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), and Instituto Nanocell.

Note

The authors declare that they have no conflicts of interest relating to this paper content.

References

1. Calhoun DA, Jones D, Textor S, Goff DC, Murphy TP, Toto RD, et al. Resistant hypertension: diagnosis, evaluation, and treatment: a scientific statement from the American Heart Association Professional Education Committee of the Council for High Blood Pressure Research. *Circulation*. 2008;117(25):e510-26.
2. Egan BM, Zhao Y, Axon RN, Brzezinski WA, Ferdinand KC. Uncontrolled and apparent treatment resistant hypertension in the United States, 1988 to 2008. *Circulation*. 2011;124(9):1046-58.
3. Sarafidis PA, Georgianos P, Bakris GL. Resistant hypertension--its identification and epidemiology. *Nature reviews Nephrology*. 2013;9(1):51-8.
4. Jung O, Gechter JL, Wunder C, Paulke A, Bartel C, Geiger H, et al. Resistant hypertension? Assessment of adherence by toxicological urine analysis. *Journal of hypertension*. 2013;31(4):766-74.
5. Muntner P, Davis BR, Cushman WC, Bangalore S, Calhoun DA, Pressel SL, et al. Treatment-resistant hypertension and the incidence of cardiovascular disease and end-stage renal disease: results from the Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT). *Hypertension*. 2014;64(5):1012-21.
6. Martins LC, Figueiredo VN, Quinaglia T, Boer-Martins L, Yugar-Toledo JC, Martin JF, et al. Characteristics of resistant hypertension: ageing, body mass index, hyperaldosteronism, cardiac hypertrophy and vascular stiffness. *Journal of human hypertension*. 2011;25(9):532-8.
7. Pedrosa RP, Drager LF, Gonzaga CC, Sousa MG, de Paula LK, Amaro AC, et al. Obstructive sleep apnea: the most common secondary cause of hypertension associated with resistant hypertension. *Hypertension*. 2011;58(5):811-7.
8. Umpierrez GE, Cantey P, Smiley D, Palacio A, Temponi D, Luster K, et al. Primary aldosteronism in diabetic subjects with resistant hypertension. *Diabetes care*. 2007;30(7):1699-703.
9. Funder JW, Carey RM, Fardella C, Gomez-Sanchez CE, Mantero F, Stowasser M, et al. Case detection, diagnosis, and treatment of patients with primary

- aldosteronism: an endocrine society clinical practice guideline. *The Journal of clinical endocrinology and metabolism*. 2008;93(9):3266-81.
10. Makris A, Seferou M, Papadopoulos DP. Resistant hypertension workup and approach to treatment. *International journal of hypertension*. 2010;2011:598694.
 11. Weber F, Anlauf M. Treatment resistant hypertension--investigation and conservative management. *Deutsches Arzteblatt international*. 2014;111(25):425-31.
 12. Mahfoud F, Himmel F, Ukena C, Schunkert H, Bohm M, Weil J. Treatment strategies for resistant arterial hypertension. *Deutsches Arzteblatt international*. 2011;108(43):725-31.
 13. Vidt DG. Contributing factors in resistant hypertension. Truly refractory disease is rarely found in a properly conducted workup. *Postgraduate medicine*. 2000;107(5):57-60, 3-4, 7-8, 70.
 14. Vongpatanasin W. Resistant hypertension: a review of diagnosis and management. *Jama*. 2014;311(21):2216-24.
 15. Taler SJ. Treatment of resistant hypertension. *Current hypertension reports*. 2005;7(5):323-9.
 16. Passarelli O, Gonçalves MS, Amodeo C. Fluxograma abordagem diagnóstica da hipertensão resistente. *Revista Brasileira de Hipertensão*. 2009;16(1):7-9.
 17. Rubin LJ, Simonneau G, Badesch D, Galie N, Humbert M, Keogh A, et al. The study of risk in pulmonary arterial hypertension. *European respiratory review : an official journal of the European Respiratory Society*. 2012;21(125):234-8.
 18. Maron BA, Leopold JA. The role of the renin-angiotensin-aldosterone system in the pathobiology of pulmonary arterial hypertension (2013 Grover Conference series). *Pulmonary circulation*. 2014;4(2):200-10.
 19. Santos RA, Ferreira AJ, Simoes ESAC. Recent advances in the angiotensin-converting enzyme 2-angiotensin(1-7)-Mas axis. *Experimental physiology*. 2008;93(5):519-27.
 20. Briet M, Schiffrin EL. Aldosterone: effects on the kidney and cardiovascular system. *Nature reviews Nephrology*. 2010;6(5):261-73.
 21. Oudit GY, Crackower MA, Backx PH, Penninger JM. The role of ACE2 in cardiovascular physiology. *Trends in cardiovascular medicine*. 2003;13(3):93-101.
 22. Gaddam KK, Nishizaka MK, Pratt-Ubunama MN, Pimenta E, Aban I, Oparil S, et al. Characterization of resistant hypertension: association between resistant hypertension, aldosterone, and persistent intravascular volume expansion. *Archives of internal medicine*. 2008;168(11):1159-64.
 23. Taler SJ, Textor SC, Augustine JE. Resistant hypertension: comparing hemodynamic management to specialist care. *Hypertension*. 2002;39(5):982-8.
 24. Boomsma F, van den Meiracker AH. Plasma A- and B-type natriuretic peptides: physiology, methodology and clinical use. *Cardiovascular research*. 2001;51(3):442-9.
 25. Freel EM, Connell JM. Mechanisms of hypertension: the expanding role of aldosterone. *Journal of the American Society of Nephrology : JASN*. 2004;15(8):1993-2001.
 26. De Martino MC, van Koetsveld PM, Pivonello R, Hofland LJ. Role of the mTOR pathway in normal and tumoral adrenal cells. *Neuroendocrinology*. 2010;92 Suppl 1:28-34.
 27. Su H, Gu Y, Li F, Wang Q, Huang B, Jin X, et al. The PI3K/AKT/mTOR signaling pathway is overactivated in primary aldosteronism. *PloS one*. 2013;8(4):e62399.
 28. Brizuela L, Rabano M, Gangoiti P, Narbona N, Macarulla JM, Trueba M, et al. Sphingosine-1-phosphate stimulates aldosterone secretion through a mechanism

- involving the PI3K/PKB and MEK/ERK 1/2 pathways. *Journal of lipid research*. 2007;48(10):2264-74.
29. Battelli C, Cho DC. mTOR inhibitors in renal cell carcinoma. *Therapy*. 2011;8(4):359-67.
 30. Brizuela L, Rabano M, Pena A, Gangoiti P, Macarulla JM, Trueba M, et al. Sphingosine 1-phosphate: a novel stimulator of aldosterone secretion. *Journal of lipid research*. 2006;47(6):1238-49.
 31. Spiegel S, Milstien S. Sphingosine-1-phosphate: an enigmatic signalling lipid. *Nature reviews*. 2003;4(5):397-407.
 32. de Almeida PWM, Lima RD, Gomes ERD, Rocha-Resende C, Roman-Campos D, Gondim ANS, et al. Functional Cross-Talk Between Aldosterone and Angiotensin-(1-7) in Ventricular Myocytes. *Hypertension*. 2013;61(2):425-+.
 33. de Almeida PW, Melo MB, Lima Rde F, Gavioli M, Santiago NM, Greco L, et al. Beneficial effects of angiotensin-(1-7) against deoxycorticosterone acetate-induced diastolic dysfunction occur independently of changes in blood pressure. *Hypertension*. 2015;66(2):389-95.
 34. de Faria AP, Demacq C, Figueiredo VN, Moraes CH, Santos RC, Sabbatini AR, et al. Hypoadiponectinemia and aldosterone excess are associated with lack of blood pressure control in subjects with resistant hypertension. *Hypertension research : official journal of the Japanese Society of Hypertension*. 2013;36(12):1067-72.
 35. Fontana V, de Faria AP, Barbaro NR, Sabbatini AR, Modolo R, Lacchini R, et al. Modulation of aldosterone levels by -344 C/T CYP11B2 polymorphism and spironolactone use in resistant hypertension. *Journal of the American Society of Hypertension : JASH*. 2014;8(3):146-51.
 36. Sampaio WO, Souza dos Santos RA, Faria-Silva R, da Mata Machado LT, Schiffrin EL, Touyz RM. Angiotensin-(1-7) through receptor Mas mediates endothelial nitric oxide synthase activation via Akt-dependent pathways. *Hypertension*. 2007;49(1):185-92.
 37. Rabelo LA, Alenina N, Bader M. ACE2-angiotensin-(1-7)-Mas axis and oxidative stress in cardiovascular disease. *Hypertension research : official journal of the Japanese Society of Hypertension*. 2011;34(2):154-60.
 38. Santos RA, Ferreira AJ, Pinheiro SV, Sampaio WO, Touyz R, Campagnole-Santos MJ. Angiotensin-(1-7) and its receptor as a potential targets for new cardiovascular drugs. *Expert opinion on investigational drugs*. 2005;14(8):1019-31.
 39. Ferreira AJ, Santos RA, Bradford CN, Mecca AP, Sumners C, Katovich MJ, et al. Therapeutic implications of the vasoprotective axis of the renin-angiotensin system in cardiovascular diseases. *Hypertension*. 2010;55(2):207-13.
 40. Gironacci MM, Longo Carbajosa NA, Goldstein J, Cerrato BD. Neuromodulatory role of angiotensin-(1-7) in the central nervous system. *Clinical science*. 2013;125(2):57-65.
 41. Gomes ER, Lara AA, Almeida PW, Guimaraes D, Resende RR, Campagnole-Santos MJ, et al. Angiotensin-(1-7) prevents cardiomyocyte pathological remodeling through a nitric oxide/guanosine 3',5'-cyclic monophosphate-dependent pathway. *Hypertension*. 2010;55(1):153-60.
 42. Passos-Silva DG, Verano-Braga T, Santos RA. Angiotensin-(1-7): beyond the cardio-renal actions. *Clinical science*. 2013;124(7):443-56.
 43. Touyz RM, Tabet F, Schiffrin EL. Redox-dependent signalling by angiotensin II and vascular remodelling in hypertension. *Clinical and experimental pharmacology & physiology*. 2003;30(11):860-6.

44. Grassi G, Ram VS. Evidence for a critical role of the sympathetic nervous system in hypertension. *Journal of the American Society of Hypertension : JASH*. 2016;10(5):457-66.
45. Bhatt DL, Kandzari DE, O'Neill WW, D'Agostino R, Flack JM, Katzen BT, et al. A controlled trial of renal denervation for resistant hypertension. *The New England journal of medicine*. 2014;370(15):1393-401.
46. Heusser K, Tank J, Brinkmann J, Menne J, Kaufeld J, Linnenweber-Held S, et al. Acute Response to Unilateral Unipolar Electrical Carotid Sinus Stimulation in Patients With Resistant Arterial Hypertension. *Hypertension*. 2016;67(3):585-91.
47. Gavioli M, Lara A, Almeida PWM, Lima AM, Damasceno DD, Rocha-Resende C, et al. Cholinergic Signaling Exerts Protective Effects in Models of Sympathetic Hyperactivity-Induced Cardiac Dysfunction. *PloS one*. 2014;9(7):e100179.
48. Paschon V, Higa GSV, Resende RR, Britto LRG, Kihara AH. Blocking of Connexin-Mediated Communication Promotes Neuroprotection during Acute Degeneration Induced by Mechanical Trauma. *PloS one*. 2012;7(9):e45449.
49. de Haro Moraes C, Figueiredo VN, de Faria AP, Barbaro NR, Sabbatini AR, Quinaglia T, et al. High-circulating leptin levels are associated with increased blood pressure in uncontrolled resistant hypertension. *Journal of human hypertension*. 2013;27(4):225-30.
50. Dardeno TA, Chou SH, Moon HS, Chamberland JP, Fiorenza CG, Mantzoros CS. Leptin in human physiology and therapeutics. *Frontiers in neuroendocrinology*. 2010;31(3):377-93.
51. Rahmouni K, Correia ML, Haynes WG, Mark AL. Obesity-associated hypertension: new insights into mechanisms. *Hypertension*. 2005;45(1):9-14.
52. Morrison CD. Leptin signaling in brain: A link between nutrition and cognition? *Biochimica et biophysica acta*. 2009;1792(5):401-8.
53. Ste Marie L, Miura GI, Marsh DJ, Yagaloff K, Palmiter RD. A metabolic defect promotes obesity in mice lacking melanocortin-4 receptors. *Proceedings of the National Academy of Sciences of the United States of America*. 2000;97(22):12339-44.
54. Kang YS. Obesity associated hypertension: new insights into mechanism. *Electrolyte & blood pressure : E & BP*. 2013;11(2):46-52.
55. Williams TM, Lisanti MP. The Caveolin genes: from cell biology to medicine. *Annals of medicine*. 2004;36(8):584-95.
56. Pojoga LH, Underwood PC, Goodarzi MO, Williams JS, Adler GK, Jeunemaitre X, et al. Variants of the caveolin-1 gene: a translational investigation linking insulin resistance and hypertension. *The Journal of clinical endocrinology and metabolism*. 2011;96(8):E1288-92.
57. Hnasko R, Lisanti MP. The biology of caveolae: lessons from caveolin knockout mice and implications for human disease. *Molecular interventions*. 2003;3(8):445-64.
58. Pojoga LH, Yao TM, Sinha S, Ross RL, Lin JC, Raffetto JD, et al. Effect of dietary sodium on vasoconstriction and eNOS-mediated vascular relaxation in caveolin-1-deficient mice. *American journal of physiology Heart and circulatory physiology*. 2008;294(3):H1258-65.
59. Cohen AW, Razani B, Wang XB, Combs TP, Williams TM, Scherer PE, et al. Caveolin-1-deficient mice show insulin resistance and defective insulin receptor protein expression in adipose tissue. *American journal of physiology Cell physiology*. 2003;285(1):C222-35.
60. Cohen AW, Combs TP, Scherer PE, Lisanti MP. Role of caveolin and caveolae in insulin signaling and diabetes. *American journal of physiology Endocrinology and metabolism*. 2003;285(6):E1151-60.

61. Karlsson M, Thorn H, Parpal S, Stralfors P, Gustavsson J. Insulin induces translocation of glucose transporter GLUT4 to plasma membrane caveolae in adipocytes. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology*. 2002;16(2):249-51.
62. Oh YS, Cho KA, Ryu SJ, Khil LY, Jun HS, Yoon JW, et al. Regulation of insulin response in skeletal muscle cell by caveolin status. *Journal of cellular biochemistry*. 2006;99(3):747-58.
63. Figueiredo VN, Yugar-Toledo JC, Martins LC, Martins LB, de Faria AP, de Haro Moraes C, et al. Vascular stiffness and endothelial dysfunction: Correlations at different levels of blood pressure. *Blood pressure*. 2012;21(1):31-8.
64. Fernandez-Patron C. Therapeutic potential of the epidermal growth factor receptor transactivation in hypertension: a convergent signaling pathway of vascular tone, oxidative stress, and hypertrophic growth downstream of vasoactive G-protein-coupled receptors? *Canadian journal of physiology and pharmacology*. 2007;85(1):97-104.
65. Hao L, Du M, Lopez-Campistrous A, Fernandez-Patron C. Agonist-induced activation of matrix metalloproteinase-7 promotes vasoconstriction through the epidermal growth factor-receptor pathway. *Circulation research*. 2004;94(1):68-76.
66. Aguiar CJ, Rocha-Franco JA, Sousa PA, Santos AK, Ladeira M, Rocha-Resende C, et al. Succinate causes pathological cardiomyocyte hypertrophy through GPR91 activation. *Cell Communication and Signaling : CCS*. 2014;12:78.
67. Nagareddy PR, MacLeod KM, McNeill JH. GPCR agonist-induced transactivation of the EGFR upregulates MLC II expression and promotes hypertension in insulin-resistant rats. *Cardiovascular research*. 2010;87(1):177-86.
68. Prenzel N, Zwick E, Daub H, Leserer M, Abraham R, Wallasch C, et al. EGF receptor transactivation by G-protein-coupled receptors requires metalloproteinase cleavage of proHB-EGF. *Nature*. 1999;402(6764):884-8.
69. Resende RR, Andrade LM, Oliveira AG, Guimaraes ES, Guatimosim S, Leite MF. Nucleoplasmic calcium signaling and cell proliferation: calcium signaling in the nucleus. *Cell Commun Signal*. 2013;11(1):14.
70. Roy A, Fields WC, Rocha-Resende C, Resende RR, Guatimosim S, Prado VF, et al. Cardiomyocyte-secreted acetylcholine is required for maintenance of homeostasis in the heart. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology*. 2013;27(12):5072-82.
71. Asakura M, Kitakaze M, Takashima S, Liao Y, Ishikura F, Yoshinaka T, et al. Cardiac hypertrophy is inhibited by antagonism of ADAM12 processing of HB-EGF: metalloproteinase inhibitors as a new therapy. *Nature medicine*. 2002;8(1):35-40.
72. Lifton RP, Gharavi AG, Geller DS. Molecular mechanisms of human hypertension. *Cell*. 2001;104(4):545-56.
73. Fujino T, Hasebe N, Fujita M, Takeuchi K, Kawabe J, Tobise K, et al. Enhanced expression of heparin-binding EGF-like growth factor and its receptor in hypertrophied left ventricle of spontaneously hypertensive rats. *Cardiovascular research*. 1998;38(2):365-74.
74. Rocha-Resende C, Roy A, Resende R, Ladeira MS, Lara A, de Moraes Gomes ER, et al. Non-neuronal cholinergic machinery present in cardiomyocytes offsets hypertrophic signals. *Journal of molecular and cellular cardiology*. 2012;53(2):206-16.
75. Arantes LA, Aguiar CJ, Amaya MJ, Figueiro NC, Andrade LM, Rocha-Resende C, et al. Nuclear inositol 1,4,5-trisphosphate is a necessary and conserved signal for the induction of both pathological and physiological cardiomyocyte hypertrophy. *Journal of molecular and cellular cardiology*. 2012;53(4):475-86.

76. Lukaszewicz KM, Lombard JH. Role of the CYP4A/20-HETE pathway in vascular dysfunction of the Dahl salt-sensitive rat. *Clinical science*. 2013;124(12):695-700.
77. Roman RJ. P-450 metabolites of arachidonic acid in the control of cardiovascular function. *Physiological reviews*. 2002;82(1):131-85.
78. Ward NC, Tsai IJ, Barden A, van Bockxmeer FM, Puddey IB, Hodgson JM, et al. A single nucleotide polymorphism in the CYP4F2 but not CYP4A11 gene is associated with increased 20-HETE excretion and blood pressure. *Hypertension*. 2008;51(5):1393-8.
79. Liu H, Zhao Y, Nie D, Shi J, Fu L, Li Y, et al. Association of a functional cytochrome P450 4F2 haplotype with urinary 20-HETE and hypertension. *Journal of the American Society of Nephrology : JASN*. 2008;19(4):714-21.
80. Drenjancevic-Peric I, Greene AS, Kunert MP, Lombard JH. Arteriolar responses to vasodilator stimuli and elevated P(O₂) in renin congenic and Dahl salt-sensitive rats. *Microcirculation*. 2004;11(8):669-77.
81. Durand MJ, Lombard JH. Introgression of the Brown Norway renin allele onto the Dahl salt-sensitive genetic background increases Cu/Zn SOD expression in cerebral arteries. *American journal of hypertension*. 2011;24(5):563-8.
82. Drenjancevic-Peric I, Lombard JH. Reduced angiotensin II and oxidative stress contribute to impaired vasodilation in Dahl salt-sensitive rats on low-salt diet. *Hypertension*. 2005;45(4):687-91.
83. Durand MJ, Moreno C, Greene AS, Lombard JH. Impaired relaxation of cerebral arteries in the absence of elevated salt intake in normotensive congenic rats carrying the Dahl salt-sensitive renin gene. *American journal of physiology Heart and circulatory physiology*. 2010;299(6):H1865-74.
84. Lenda DM, Boegehold MA. Effect of a high salt diet on microvascular antioxidant enzymes. *Journal of vascular research*. 2002;39(1):41-50.
85. Ha M, Kim VN. Regulation of microRNA biogenesis. *Nature reviews*. 2014;15(8):509-24.
86. Ambros V, Bartel B, Bartel DP, Burge CB, Carrington JC, Chen X, et al. A uniform system for microRNA annotation. *Rna*. 2003;9(3):277-9.
87. Zeng Y, Yi R, Cullen BR. MicroRNAs and small interfering RNAs can inhibit mRNA expression by similar mechanisms. *Proceedings of the National Academy of Sciences of the United States of America*. 2003;100(17):9779-84.
88. Lee Y, Ahn C, Han J, Choi H, Kim J, Yim J, et al. The nuclear RNase III Drosha initiates microRNA processing. *Nature*. 2003;425(6956):415-9.
89. Basyuk E, Suavet F, Doglio A, Bordonne R, Bertrand E. Human let-7 stem-loop precursors harbor features of RNase III cleavage products. *Nucleic acids research*. 2003;31(22):6593-7.
90. Lund E, Guttinger S, Calado A, Dahlberg JE, Kutay U. Nuclear export of microRNA precursors. *Science*. 2004;303(5654):95-8.
91. Bernstein E, Caudy AA, Hammond SM, Hannon GJ. Role for a bidentate ribonuclease in the initiation step of RNA interference. *Nature*. 2001;409(6818):363-6.
92. Hutvagner G, Zamore PD. A microRNA in a multiple-turnover RNAi enzyme complex. *Science*. 2002;297(5589):2056-60.
93. Schwarz DS, Hutvagner G, Du T, Xu Z, Aronin N, Zamore PD. Asymmetry in the assembly of the RNAi enzyme complex. *Cell*. 2003;115(2):199-208.
94. Blackshaw S, Harpavat S, Trimarchi J, Cai L, Huang H, Kuo WP, et al. Genomic analysis of mouse retinal development. *PLoS Biol*. 2004;2(9):E247.

95. Chen X, Ba Y, Ma L, Cai X, Yin Y, Wang K, et al. Characterization of microRNAs in serum: a novel class of biomarkers for diagnosis of cancer and other diseases. *Cell research*. 2008;18(10):997-1006.
96. Higa GS, de Sousa E, Walter LT, Kinjo ER, Resende RR, Kihara AH. MicroRNAs in neuronal communication. *Molecular neurobiology*. 2014;49(3):1309-26.
97. Pinto MC, Tonelli FM, Vieira AL, Kihara AH, Ulrich H, Resende RR. Studying complex system: calcium oscillations as attractor of cell differentiation. *Integrative biology : quantitative biosciences from nano to macro*. 2016;8(2):130-48.
98. Vasconcellos R, Alvarenga EC, Parreira RC, Lima SS, Resende RR. Exploring the cell signalling in hepatocyte differentiation. *Cellular signalling*. 2016;28(11):1773-88.
99. van Rooij E, Sutherland LB, Liu N, Williams AH, McAnally J, Gerard RD, et al. A signature pattern of stress-responsive microRNAs that can evoke cardiac hypertrophy and heart failure. *Proceedings of the National Academy of Sciences of the United States of America*. 2006;103(48):18255-60.
100. Hoekstra M, van der Lans CA, Halvorsen B, Gullestad L, Kuiper J, Aukrust P, et al. The peripheral blood mononuclear cell microRNA signature of coronary artery disease. *Biochemical and biophysical research communications*. 2010;394(3):792-7.
101. Batkai S, Thum T. MicroRNAs in hypertension: mechanisms and therapeutic targets. *Current hypertension reports*. 2012;14(1):79-87.
102. Mitchell PS, Parkin RK, Kroh EM, Fritz BR, Wyman SK, Pogosova-Agadjanyan EL, et al. Circulating microRNAs as stable blood-based markers for cancer detection. *Proceedings of the National Academy of Sciences of the United States of America*. 2008;105(30):10513-8.
103. Li C, Pei F, Zhu X, Duan DD, Zeng C. Circulating microRNAs as novel and sensitive biomarkers of acute myocardial Infarction. *Clinical biochemistry*. 2012;45(10-11):727-32.
104. Marques FZ, Campain AE, Tomaszewski M, Zukowska-Szczechowska E, Yang YH, Charchar FJ, et al. Gene expression profiling reveals renin mRNA overexpression in human hypertensive kidneys and a role for microRNAs. *Hypertension*. 2011;58(6):1093-8.
105. Marques FZ, Romaine SP, Denniff M, Eales J, Dormer J, Garrelds IM, et al. Signatures of miR-181a on renal transcriptome and blood pressure. *Molecular medicine*. 2015.
106. Sansom SE, Nuovo GJ, Martin MM, Kotha SR, Parinandi NL, Elton TS. miR-802 regulates human angiotensin II type 1 receptor expression in intestinal epithelial C2BBel cells. *American journal of physiology Gastrointestinal and liver physiology*. 2010;299(3):G632-42.
107. MicroRNA-155 regulates human angiotensin II type 1 receptor expression in fibroblasts. *The Journal of biological chemistry*. 2013;288(6):4226.
108. Zheng L, Xu CC, Chen WD, Shen WL, Ruan CC, Zhu LM, et al. MicroRNA-155 regulates angiotensin II type 1 receptor expression and phenotypic differentiation in vascular adventitial fibroblasts. *Biochemical and biophysical research communications*. 2010;400(4):483-8.
109. Li S, Zhu J, Zhang W, Chen Y, Zhang K, Popescu LM, et al. Signature microRNA expression profile of essential hypertension and its novel link to human cytomegalovirus infection. *Circulation*. 2011;124(2):175-84.
110. Kemp JR, Unal H, Desnoyer R, Yue H, Bhatnagar A, Karnik SS. Angiotensin II-regulated microRNA 483-3p directly targets multiple components of the renin-angiotensin system. *Journal of molecular and cellular cardiology*. 2014;75:25-39.

111. Eskildsen TV, Jeppesen PL, Schneider M, Nossent AY, Sandberg MB, Hansen PB, et al. Angiotensin II regulates microRNA-132/-212 in hypertensive rats and humans. *International journal of molecular sciences*. 2013;14(6):11190-207.
112. Nossent AY, Hansen JL, Doggen C, Quax PH, Sheikh SP, Rosendaal FR. SNPs in microRNA binding sites in 3'-UTRs of RAAS genes influence arterial blood pressure and risk of myocardial infarction. *American journal of hypertension*. 2011;24(9):999-1006.
113. Maatta K, Kunnas T, Nikkari ST. Contribution of SLC7A1 genetic variant to hypertension, the TAMRISK study. *BMC medical genetics*. 2013;14:69.
114. Yang Z, Venardos K, Jones E, Morris BJ, Chin-Dusting J, Kaye DM. Identification of a novel polymorphism in the 3'UTR of the L-arginine transporter gene SLC7A1: contribution to hypertension and endothelial dysfunction. *Circulation*. 2007;115(10):1269-74.
115. Yang Z, Kaye DM. Mechanistic insights into the link between a polymorphism of the 3'UTR of the SLC7A1 gene and hypertension. *Human mutation*. 2009;30(3):328-33.
116. Robertson S, MacKenzie SM, Alvarez-Madrazo S, Diver LA, Lin J, Stewart PM, et al. MicroRNA-24 is a novel regulator of aldosterone and cortisol production in the human adrenal cortex. *Hypertension*. 2013;62(3):572-8.
117. Fiedler J, Jazbutyte V, Kirchmaier BC, Gupta SK, Lorenzen J, Hartmann D, et al. MicroRNA-24 regulates vascularity after myocardial infarction. *Circulation*. 2011;124(6):720-30.
118. Meloni M, Marchetti M, Garner K, Littlejohns B, Sala-Newby G, Xenophontos N, et al. Local inhibition of microRNA-24 improves reparative angiogenesis and left ventricle remodeling and function in mice with myocardial infarction. *Molecular therapy : the journal of the American Society of Gene Therapy*. 2013;21(7):1390-402.
119. Sober S, Laan M, Annilo T. MicroRNAs miR-124 and miR-135a are potential regulators of the mineralocorticoid receptor gene (NR3C2) expression. *Biochemical and biophysical research communications*. 2010;391(1):727-32.
120. Mopidevi B, Ponnala M, Kumar A. Human angiotensinogen +11525 C/A polymorphism modulates its gene expression through microRNA binding. *Physiological genomics*. 2013;45(19):901-6.
121. Elia L, Quintavalle M, Zhang J, Contu R, Cossu L, Latronico MV, et al. The knockout of miR-143 and -145 alters smooth muscle cell maintenance and vascular homeostasis in mice: correlates with human disease. *Cell death and differentiation*. 2009;16(12):1590-8.
122. Albinsson S, Skoura A, Yu J, DiLorenzo A, Fernandez-Hernando C, Offermanns S, et al. Smooth muscle miRNAs are critical for post-natal regulation of blood pressure and vascular function. *PloS one*. 2011;6(4):e18869.
123. Boettger T, Beetz N, Kostin S, Schneider J, Kruger M, Hein L, et al. Acquisition of the contractile phenotype by murine arterial smooth muscle cells depends on the Mir143/145 gene cluster. *The Journal of clinical investigation*. 2009;119(9):2634-47.
124. Xin M, Small EM, Sutherland LB, Qi X, McAnally J, Plato CF, et al. MicroRNAs miR-143 and miR-145 modulate cytoskeletal dynamics and responsiveness of smooth muscle cells to injury. *Genes Dev*. 2009;23(18):2166-78.
125. Kontaraki JE, Marketou ME, Zacharis EA, Parthenakis FI, Vardas PE. MicroRNA-9 and microRNA-126 expression levels in patients with essential hypertension: potential markers of target-organ damage. *Journal of the American Society of Hypertension : JASH*. 2014;8(6):368-75.

126. Wang K, Long B, Zhou J, Li PF. miR-9 and NFATc3 regulate myocardin in cardiac hypertrophy. *The Journal of biological chemistry*. 2010;285(16):11903-12.
127. Wang S, Aurora AB, Johnson BA, Qi X, McAnally J, Hill JA, et al. The endothelial-specific microRNA miR-126 governs vascular integrity and angiogenesis. *Developmental cell*. 2008;15(2):261-71.
128. Nicoli S, Standley C, Walker P, Hurlstone A, Fogarty KE, Lawson ND. MicroRNA-mediated integration of haemodynamics and Vegf signalling during angiogenesis. *Nature*. 2010;464(7292):1196-200.
129. Harris TA, Yamakuchi M, Ferlito M, Mendell JT, Lowenstein CJ. MicroRNA-126 regulates endothelial expression of vascular cell adhesion molecule 1. *Proceedings of the National Academy of Sciences of the United States of America*. 2008;105(5):1516-21.
130. Sekar D, Shilpa BR, Das AJ. Relevance of microRNA 21 in Different Types of Hypertension. *Current hypertension reports*. 2017;19(7):57.
131. Sabatel C, Malvaux L, Bovy N, Deroanne C, Lambert V, Gonzalez ML, et al. MicroRNA-21 exhibits antiangiogenic function by targeting RhoB expression in endothelial cells. *PloS one*. 2011;6(2):e16979.
132. Fleissner F, Jazbutyte V, Fiedler J, Gupta SK, Yin X, Xu Q, et al. Short communication: asymmetric dimethylarginine impairs angiogenic progenitor cell function in patients with coronary artery disease through a microRNA-21-dependent mechanism. *Circulation research*. 2010;107(1):138-43.
133. Jazbutyte V, Thum T. MicroRNA-21: from cancer to cardiovascular disease. *Current drug targets*. 2010;11(8):926-35.
134. Ji R, Cheng Y, Yue J, Yang J, Liu X, Chen H, et al. MicroRNA expression signature and antisense-mediated depletion reveal an essential role of MicroRNA in vascular neointimal lesion formation. *Circulation research*. 2007;100(11):1579-88.
135. Liu X, Cheng Y, Zhang S, Lin Y, Yang J, Zhang C. A necessary role of miR-221 and miR-222 in vascular smooth muscle cell proliferation and neointimal hyperplasia. *Circulation research*. 2009;104(4):476-87.
136. Davis BN, Hilyard AC, Nguyen PH, Lagna G, Hata A. Induction of microRNA-221 by platelet-derived growth factor signaling is critical for modulation of vascular smooth muscle phenotype. *The Journal of biological chemistry*. 2009;284(6):3728-38.
137. Menghini R, Casagrande V, Cardellini M, Martelli E, Terrinoni A, Amati F, et al. MicroRNA 217 modulates endothelial cell senescence via silent information regulator 1. *Circulation*. 2009;120(15):1524-32.
138. Liu Y, Taylor NE, Lu L, Usa K, Cowley AW, Jr., Ferreri NR, et al. Renal medullary microRNAs in Dahl salt-sensitive rats: miR-29b regulates several collagens and related genes. *Hypertension*. 2010;55(4):974-82.
139. Zhang Y, Huang XR, Wei LH, Chung AC, Yu CM, Lan HY. miR-29b as a therapeutic agent for angiotensin II-induced cardiac fibrosis by targeting TGF-beta/Smad3 signaling. *Molecular therapy : the journal of the American Society of Gene Therapy*. 2014;22(5):974-85.
140. Jackson KL, Marques FZ, Watson AM, Palma-Rigo K, Nguyen-Huu TP, Morris BJ, et al. A novel interaction between sympathetic overactivity and aberrant regulation of renin by miR-181a in BPH/2J genetically hypertensive mice. *Hypertension*. 2013;62(4):775-81.
141. Head GA, Gueguen C, Marques FZ, Jackson KL, Eikelis N, Stevenson ER, et al. 6b.01: Effect of Renal Denervation on Blood Pressure and Microrna 181a in Hypertensive Schlager Mice. *Journal of hypertension*. 2015;33 Suppl 1:e76.

142. Dorr O, Liebetrau C, Mollmann H, Gaede L, Troidl C, Lankes S, et al. Effect of Renal Sympathetic Denervation on Specific MicroRNAs as an Indicator of Reverse Remodeling Processes in Hypertensive Heart Disease. *Journal of clinical hypertension*. 2016;18(6):497-502.
143. Gu N, You L, Shi C, Yang L, Pang L, Cui X, et al. Expression of miR-199a-3p in human adipocytes is regulated by free fatty acids and adipokines. *Molecular medicine reports*. 2016;14(2):1180-6.
144. Santovito D, De Nardis V, Marcantonio P, Mandolini C, Paganelli C, Vitale E, et al. Plasma exosome microRNA profiling unravels a new potential modulator of adiponectin pathway in diabetes: effect of glycemic control. *The Journal of clinical endocrinology and metabolism*. 2014;99(9):E1681-5.
145. Zampetaki A, Kiechl S, Drozdov I, Willeit P, Mayr U, Prokopi M, et al. Plasma microRNA profiling reveals loss of endothelial miR-126 and other microRNAs in type 2 diabetes. *Circulation research*. 2010;107(6):810-7.
146. Pulakat L, Aroor AR, Gul R, Sowers JR. Cardiac insulin resistance and microRNA modulators. *Experimental diabetes research*. 2012;2012:654904.
147. Lee Y, Kim M, Han J, Yeom KH, Lee S, Baek SH, et al. MicroRNA genes are transcribed by RNA polymerase II. *The EMBO journal*. 2004;23(20):4051-60.
148. Guo CJ, Pan Q, Xiong H, Qiao YQ, Bian ZL, Zhong W, et al. Therapeutic potential of microRNA: a new target to treat intrahepatic portal hypertension? *BioMed research international*. 2014;2014:797898.
149. Lozano L, Tovar JL, Sampol G, Romero O, Jurado MJ, Segarra A, et al. Continuous positive airway pressure treatment in sleep apnea patients with resistant hypertension: a randomized, controlled trial. *Journal of hypertension*. 2010;28(10):2161-8.
150. Martinez-Garcia MA, Capote F, Campos-Rodriguez F, Lloberes P, Diaz de Atauri MJ, Somoza M, et al. Effect of CPAP on blood pressure in patients with obstructive sleep apnea and resistant hypertension: the HIPARCO randomized clinical trial. *Jama*. 2013;310(22):2407-15.
151. Calhoun DA, Nishizaka MK, Zaman MA, Thakkar RB, Weissmann P. Hyperaldosteronism among black and white subjects with resistant hypertension. *Hypertension*. 2002;40(6):892-6.
152. Sang X, Jiang Y, Wang W, Yan L, Zhao J, Peng Y, et al. Prevalence of and risk factors for primary aldosteronism among patients with resistant hypertension in China. *Journal of hypertension*. 2013;31(7):1465-71; discussion 71-2.
153. Parthasarathy HK, Menard J, White WB, Young WF, Jr., Williams GH, Williams B, et al. A double-blind, randomized study comparing the antihypertensive effect of eplerenone and spironolactone in patients with hypertension and evidence of primary aldosteronism. *Journal of hypertension*. 2011;29(5):980-90.
154. Rossi GP, Cesari M, Cuspidi C, Maiolino G, Cicala MV, Bisogni V, et al. Long-term control of arterial hypertension and regression of left ventricular hypertrophy with treatment of primary aldosteronism. *Hypertension*. 2013;62(1):62-9.
155. Benjamin MM, Fazel P, Filardo G, Choi JW, Stoler RC. Prevalence of and risk factors of renal artery stenosis in patients with resistant hypertension. *The American journal of cardiology*. 2014;113(4):687-90.
156. Cooper CJ, Murphy TP, Cutlip DE, Jamerson K, Henrich W, Reid DM, et al. Stenting and medical therapy for atherosclerotic renal-artery stenosis. *The New England journal of medicine*. 2014;370(1):13-22.

157. Trinquart L, Mounier-Vehier C, Sapoval M, Gagnon N, Plouin PF. Efficacy of revascularization for renal artery stenosis caused by fibromuscular dysplasia: a systematic review and meta-analysis. *Hypertension*. 2010;56(3):525-32.
158. Shimbo D, Levitan EB, Booth JN, 3rd, Calhoun DA, Judd SE, Lackland DT, et al. The contributions of unhealthy lifestyle factors to apparent resistant hypertension: findings from the Reasons for Geographic And Racial Differences in Stroke (REGARDS) study. *Journal of hypertension*. 2013;31(2):370-6.
159. de Souza F, Muxfeldt E, Fiszman R, Salles G. Efficacy of spironolactone therapy in patients with true resistant hypertension. *Hypertension*. 2010;55(1):147-52.

Figure Legends

FIGURE 1: Recommendations for Diagnostics of RH. ABPM: Ambulatory Blood Pressure Monitoring; BP: Blood Pressure; RH: Resistant Hypertension; WCH: White-Coat Hypertension. Modified Calhoun (1) and Passarelli, Gonçalves (16)

FIGURE 2: Hemodynamic control of Renin-Angiotensin-Aldosterone System. ACE: Angiotensin Converting Enzyme; Modified Maron and Leopold (18)

FIGURE 3: Signaling pathway Na^+/k^+ -ATPase activity and Aldosterone.

FIGURE 4: Signalling pathway involved in Primary Aldosteronism. S1P – Sphingosine-1-phosphate; HB-EGF – Heparin binding-Epidermal growth factor; EGFR – Epidermal growth factor receptor; GPCR – G protein coupled receptor; PLD – Phospholipase; PKC – Protein kinase C; PI3K –Phosphoinositide 3 kinase; PIP2 – Phosphatidylinositol 4,5-biphosphate; PIP3 – Phosphatidylinositol (3,4,5)-triphosphate; AKT – Protein kinase B; mTOR – Mammalian target of rapamycin; MEK – mitogen-activated protein kinase; ERK – extracellular regulated kinase.

FIGURE 5: Signaling pathway of the SRAA axis. The figure show the regulate processes for the Ang-(1-7)/Mas in different signaling pathways, compare to the opposes RAAS classical axes of this system, the ACE/AngII/AT₁R. ANG-(1-7) – Angiotensin-(1-7); GPCR – G protein coupled receptor; PLC – Phospholipase C; DAG – Diacylglycerol; IP3 – Inositol triphosphate; PKC – Protein kinase C; PI3K – Phosphoinositide 3 kinase; PIP2 – Phosphatidylinositol 4,5-biphosphate; PIP3 –

Phosphatidylinositol (3,4,5)-triphosphate; AKT – Protein kinase B; NO – Nitric Oxide; COX-2 – Cyclooxygenase-2; PG – Prostaglandin; ERK – extracellular regulated kinase.

FIGURE 6: Signaling Leptin pathway. The figure outlines the relationship between Leptin and mechanisms involved in the neurons signaling as stimulates the sympathetic nervous system, processes involved in pathogenesis of hypertension link to obesity. SNS: Sympathetic Nervous System; RAS: Renin-Angiotensin System; FFA: Free Fatty Acids; NO: Nitric Oxide. Modified Rahmouni (51)

FIGURE 7: CAV1 Signaling Pathway. The stimulate of tyrosine kinase insulin receptor CAV1 signaling pathway lead activate PI3K resulting in translocation of the vesicle and exhibition of the GLUT4 in cell membrane. CAV1 – caveolin-1; GM3 – ganglioside; IRS-1 – insulin receptor substrate 1; PI3K –Phosphoinositide 3 kinase; PDK - Phosphoinositide-dependent kinase-1; AKT - Akt/protein kinase B; PIP2 – Phosphatidylinositol 4,5-biphosphate; PIP3 – Phosphatidylinositol (3,4,5)-triphosphate; GLUT4 – glucose transporter 4. AS160 - Akt substrate of 160 kDa.

FIGURE 8: Signaling pathways involved in the development and progression of RH. The figure illustrates the existing relationship between cell signaling mechanisms involved in the pathogenesis of hypertension. AGII – Angiotensin II; ET1 – Endothelin 1; S1P – Sphingosine-1-phosphate; NOX1 – NADPH oxidase 1; ROS – Reactive oxygen species; MMP – Matrix Metalloproteinase; HB-EGF – Heparin binding-Epidermal growth factor; ANG-(1-7) – Angiotensin-(1-7); GPCR – G protein coupled receptor;; PLA2 – Phospholipase A2; PLC – Phospholipase C; AA – Arachidonic Acid; CYP4A – Cytochrome P450-4A; MAPK – Mitogen-activated protein kinase; NO – Nitric oxide; PLD – Phospholipase; DAG – Diacylglycerol; IP3 – Inositol triphosphate; PKC – Protein kinase C; PI3K –Phosphoinositide 3 kinase; PIP2 – Phosphatidylinositol 4,5-biphosphate; PIP3 – Phosphatidylinositol (3,4,5)-triphosphate; AKT – Protein kinase B; mTOR – Mammalian target of rapamycin; COX-2 – Cyclooxygenase-2; PG – Prostaglandin; MEK – mitogen-activated protein kinase; ERK – extracellular regulated kinase; IR – insulin receptor; CAV1 – caveolin-1.

FIGURE 9: EGFR Signaling Pathway in vascular smooth muscle. The agonist action of the GPCR receptors stimulates NOX1 phatway lead to the increase of the oxidative stress NOX1which the activating the signaling pathway of the EGFR. – NADPH oxidase 1; ROS – Reactive oxygen species; MMP – Matrix Metalloproteinase; HB-EGF – Heparin binding-Epidermal growth factor; EGFR – Epidermal growth factor receptor; GPCR – G protein coupled receptor; PLC – Phospholipase C; DAG – Diacylglycerol; IP3 – Inositol triphosphate; PKC – Protein kinase C; PI3K – Phosphoinositide 3 kinase; PIP2 – Phosphatidylinositol 4,5-biphosphate; PIP3 – Phosphatidylinositol (3,4,5)-triphosphate; AKT – Protein kinase B; mTOR – Mammalian target of rapamycin.

FIGURE 10: Ang II stimulates signaling pathway through 20-HETE. The stimulation of GPCR by AngII agonist activates PLA2 which starts the signaling intracellular pathway leading to activation of PKC as release calcium influx. ROS – Reactive oxygen species; GPCR – G protein coupled receptor; PLA2 – Phospholipase A2; CYP4A – Cytochrome P450-4A; COX – Cyclooxygenase; LO – Lipoxygenase ; LTs – Leukotrienes; 20-HETE – 20-hydroxyeicosatetraenoic acid; PGE2 – Prostaglandin E2; PGI2 – Prostacyclin; PGH2 – Prostaglandin H2; TXA2 – Tromboxane A2; MAPK – Mitogen-activated protein kinase; NO – Nitric oxide;

FIGURE 11: Biogenesis of miRNAs in mammals. 1. miRNAs are transcribed in the nucleus by RNA polymerase II from long precursors. 2. In the nucleus they are processed by a complex formed by Drosha and its cofactor DGCR8. 3. The pre-miRNAs are transported to the cytoplasm by the Exportin-5 receptor. 4. In the cytoplasm, the pre-miRNAs are processed by Dicer resulting in the formation of miRNA:miRNA duplex. 5. The miRNA:miRNA duplex is processed by helicase. 6. Association with RISC complex. 7. Repression or inhibition of mRNA.

FIGURE 12. Scheme of miRNA pathway in Hypertension. Scheme of miRNA pathway in Hypertension. Major microRNAs involved in the hypertension pathogenesis. In red, those that act on muscle cells; In green, those present on cardiomyocytes; In blue, those that are expressed in endothelial cells; In brown, those that act on fibroblasts; In purple, present in renal cells; in yellow, those that are expressed in adipocytes; In black, the present in plasma. NO – Nitric Oxide; ERK – extracellular regulated kinase, GATA2 - Endothelial transcription factor GATA2; PAK4 - Serine/threonine-protein kinase PAK4; CYP11B1 - 11 β -hydroxylase CYP11B2 - aldosterone synthase; Sprouty-related protein (Sprd-1); Vascular Smooth Muscle Cells (VSMCs); C-X-C motif ligand 12 - CXCL12; nuclear receptor of subfamily 3 group C member 2 - NR3C2; monocyte chemoattractant protein-1 -MCP-1; Angiotensin type 2 receptor - AT2R; Cyclin-dependent kinase inhibitor 1B - p27, Kip1; Cyclin-dependent kinase inhibitor 1C -p57, Kip2; Vascular Endothelial Growth Factor (VEGF); Vascular Cell Adhesion Molecule 1 (VCAM-1); Ras Homolog Family Member B (RhoB), Phosphoinositide-3-kinase regulatory subunit 2 (PIK3R2); Phosphatase and tensin homolog (PTEN); Protein sprouty homolog 2 (Sprouty-2); Angiotensin-Converting Enzyme (ACE); ; Silent information regulator 1 (SirT1); Angiotensin II Receptor (AT1R); Angiotensinogen (AGT); arginine-vasopressin receptor (AVPR1A); bradykinin β 2 receptor (BDKRB2); thromboxane A2 receptor (TBXA2R); B-cell lymphoma 2 (Bcl-2)

TABLE 1: Forms of Secondary Hypertension Associated with RH. Modified Vongpatanasin (14)

TABLE 2: miRNAs and hypertension

FIGURE 1: Recommendations for Diagnostics of RH. ABPM: Ambulatory Blood Pressure Monitoring; BP: Blood Pressure; RH: Resistant Hypertension; WCH: White-Coat Hypertension. Modified Calhoun (1) and Passarelli, Gonçalves (16)

FIGURE 2: Hemodynamic control of Renin-Angiotensin-Aldosterone System. ACE: Angiotensin Converting Enzyme; Modified Maron and Leopold (18)

FIGURE 3: Signaling pathway Na^+/k^+ -ATPase activity and Aldosterone.

FIGURE 4: Signalling pathway involved in Primary Aldosteronism. S1P – Sphingosine-1-phosphate; HB-EGF – Heparin binding-Epidermal growth factor; EGFR – Epidermal growth factor receptor; GPCR – G protein coupled receptor; PLD – Phospholipase; PKC – Protein kinase C; PI3K – Phosphoinositide 3 kinase; PIP2 – Phosphatidylinositol 4,5-bisphosphate; PIP3 – Phosphatidylinositol (3,4,5)-triphosphate; AKT – Protein kinase B; mTOR – Mammalian target of rapamycin; MEK – mitogen-activated protein kinase; ERK – extracellular regulated kinase.

FIGURE 5: Signaling pathway of the SRAA axis. The figure show the regulate processes for the Ang-(1-7)/Mas in different signaling pathways, compere to the opposes RAAS classical axes of this system, the ACE/AngII/AT₁R. ANG-(1-7) – Angiotensin-(1-7); GPCR – G protein coupled receptor; PLC – Phospholipase C; DAG – Diacylglycerol; IP3 – Inositol triphosphate; PKC – Protein kinase C; PI3K – Phosphoinositide 3 kinase; PIP2 – Phosphatidylinositol 4,5-bisphosphate; PIP3 – Phosphatidylinositol (3,4,5)-triphosphate; AKT – Protein kinase B; NO – Nitric Oxide; COX-2 – Cyclooxygenase-2; PG – Prostaglandin; ERK – extracellular regulated kinase.

FIGURE 6: Signaling Leptin pathway. The figure outlines the relationship between Leptin and mechanisms involved in the neurons signaling as stimulates the sympathetic nervous system, processes involved in pathogenesis of hypertension link to obesity. SNS: Sympathetic Nervous System; RAS: Renin-Angiotensin System; FFA: Free Fatty Acids; NO: Nitric Oxide. Modified Rahmouni (51)

FIGURE 7: CAV1 Signaling Pathway. The stimulate of tyrosine kinase insulin receptor CAV1 signaling pathway lead activate PI3K resulting in translocation of the vesicle and exhibition of the GLUT4 in cell membrane. CAV1 – caveolin-1; GM3 – ganglioside; IRS-1 – insulin receptor substrate 1; PI3K – Phosphoinositide 3 kinase; PDK – Phosphoinositide-dependent kinase-1; AKT – Akt/protein kinase B; PIP2 – Phosphatidylinositol 4,5-bisphosphate; PIP3 – Phosphatidylinositol (3,4,5)-triphosphate; GLUT4 – glucose transporter 4. AS160 – Akt substrate of 160 kDa.

FIGURE 8: Signaling pathways involved in the development and progression of RH. The figure illustrates the existing relationship between cell signaling mechanisms involved in the pathogenesis of hypertension. AGII – Angiotensin II; ET1 – Endothelin 1; S1P – Sphingosine-1-phosphate; NOX1 – NADPH oxidase 1; ROS – Reactive oxygen species; MMP – Matrix Metalloproteinase; HB-EGF – Heparin binding-Epidermal growth factor; ANG-(1-7) – Angiotensin-(1-7); GPCR – G protein coupled receptor;; PLA2 – Phospholipase A2; PLC – Phospholipase C; AA – Arachidonic Acid; CYP4A – Cytochrome P450-4A; MAPK – Mitogen-activated protein kinase; NO – Nitric oxide; PLD – Phospholipase; DAG – Diacylglycerol; IP3 – Inositol triphosphate; PKC – Protein kinase C; PI3K –Phosphoinositide 3 kinase; PIP2 – Phosphatidylinositol 4,5-bisphosphate; PIP3 – Phosphatidylinositol (3,4,5)-triphosphate; AKT – Protein kinase B; mTOR – Mammalian target of rapamycin; COX-2 – Cyclooxygenase-2; PG – Prostaglandin; MEK – mitogen-activated protein kinase; ERK – extracellular regulated kinase; IR – insulin receptor; *CAVI* – caveolin-1.

FIGURE 9: EGFR Signaling Pathway in vascular smooth muscle. The agonist action of the GPCR receptors stimulates NOX1 pathway lead to the increase of the oxidative stress NOX1 which the activating the signaling pathway of the EGFR. – NADPH oxidase 1; ROS – Reactive oxygen species; MMP – Matrix Metalloproteinase; HB-EGF – Heparin binding-Epidermal growth factor; EGFR – Epidermal growth factor receptor; GPCR – G protein coupled receptor; PLC – Phospholipase C; DAG – Diacylglycerol; IP3 – Inositol triphosphate; PKC – Protein kinase C; PI3K – Phosphoinositide 3 kinase; PIP2 – Phosphatidylinositol 4,5-bisphosphate; PIP3 – Phosphatidylinositol (3,4,5)-triphosphate; AKT – Protein kinase B; mTOR – Mammalian target of rapamycin.

FIGURE 10: Ang II stimulates signaling pathway through 20-HETE. The stimulate of GPCR by AngII agonist activate PLA2 which shoots the signaling intracellular pathway leading activation of PKC as release calcium influx. ROS – Reactive oxygen species; GPCR – G protein coupled receptor; PLA2 – Phospholipase A2; CYP4A – Cytochrome P450-4A; COX – Cyclooxygenase; LO – Lipoxygenase ; LTs – Leukotrienes; 20-HETE – 20-hydroxyeicosatetraenoic acid; PGE2 – Prostaglandin E2; PGI2 – Prostacyclin; PGH2 – Prostaglandin H2; TXA2 – Tromboxane A2; MAPK – Mitogen-activated protein kinase; NO – Nitric oxide;

FIGURE 11: Biogenesis of miRNAs in mammals. 1. miRNAs are transcribed in the nucleus by RNA polymerase II from long precursors. 2. In the nucleus they are processed by a complex formed by Drosha and its cofactor DGCR8. 3. The pre-miRNAs are transported to the cytoplasm by the Exportin-5 receptor. 4. In the cytoplasm, the pre-miRNAs are processed by Dicer resulting in the formation of miRNA:miRNA duplex 5. The miRNA:miRNA duplex is processed by helicase 6. Association with RISC complex. 7. Repression or inhibition of mRNA.

FIGURE 12. Scheme of miRNA pathway in Hypertension. Scheme of miRNA pathway in Hypertension. Major microRNAs involved in the hypertension pathogenesis. In red, those that act on muscle cells; In green, those presents on cardiomyocytes; In blue, those that are expressed in endothelial cells; In brown, those that act on fibroblasts; In purple, present in renal cells; in yellow, those that are expressed in adipocytes; In black, the present in plasma. NO – Nitric Oxide; ERK – extracellular regulated kinase, GATA2 - Endothelial transcription factor GATA2; PAK4 - Serine/threonine-protein kinase PAK4; CYP11B1 - 11 β -hydroxylase CYP11B2 - aldosterone synthase; Sprouty-related protein (Spred-1); Vascular Smooth Muscle Cells (VSMCs); C-X-C motif ligand 12 - CXCL12; nuclear receptor of subfamily 3 group C member 2 - NR3C2; monocyte chemoattractant protein-1 -MCP-1; Angiotensin type 2 receptor - AT2R; Cyclin-dependent kinase inhibitor 1B - p27, Kip1; Cyclin-dependent kinase inhibitor 1C -p57, Kip2; Vascular Endothelial Growth Factor (VEGF); Vascular Cell Adhesion Molecule 1 (VCAM-1); Ras Homolog Family Member B (RhoB), Phosphoinositide-3-kinase regulatory subunit 2 (PIK3R2); Phosphatase and tensin homolog (PTEN); Protein sprouty homolog 2 (Sprouty-2); Angiotensin-Converting Enzyme (ACE); ; Silent information regulator 1 (SirT1); Angiotensin II Receptor (AT1R); Angiotensinogen (AGT); arginine-vasopressin receptor (AVPR1A); bradykinin β 2 receptor (BDKRB2); thromboxane A2 receptor (TBXA2R); B-cell lymphoma 2 (Bcl-2)

TABLE 1: Forms of Secondary Hypertension Associated with RH. Modified Vongpatanasin (14)

Conditions	Obstructive Sleep Apnea	Primary Aldosteronism	Renal Artery Stenosis	Renal Parenchyma Disease	Use of drugs and alcohol	Thyroid Disorders
Diagnostic Tests	Polysomnography	Serum aldosterone, plasma renin activity	Duplex Doppler ultrasonography, Computed tomographic angiography, or magnetic resonance angiography	Serum creatinine	History taking	Thyrotropin, free thyroxine
Treatment	Continuous positive airway Pressure	Spironolactone, eplerenone, or surgical resection of tumor in unilateral aldosterone-producing Adenoma	Renal revascularization in selected patients	Correction of underlying causes if possible	Discontinuation of offending agents	According to underlying disorders
Prevalence in RH (%)	60-70	7-20	2-24	1-2	2-4	<1
References	(7, 149, 150)	(8, 9, 151-154)	(7, 155-157)	(1, 7)	(1, 7, 158)	(1, 7)

Table 2. miRNAs related to hypertension


MiRNA	Cells Type	Target	References
miR-126	Endothelial apoptotic bodies Endothelial cells Endothelial apoptotic bodies Plasma Zebrafish embryos Endothelial cells Peripheral blood mononuclear cells Mouse Endothelial cells	VEGF VCAM-1 SPRED-1 and PIK3R2	(127-129, 138, 145)
miR-21	Endothelial cells Angiogenic progenitor cells Rat vascular smooth muscle cells Murine hearts Vascular smooth muscle cells	RhoB Superoxide dismutase 2 Sprouty-2 PTEN Bcl-2	(130-134)
miR-143 and miR-145	Mice/Rat/Mouse VSMCs	ACE Modulators of SRF activity	(121-124)
miR-221 and miR-222	Rat VSMCs	p27 (Kip1) p57 (Kip2) c-Kit	(135-136)
miR-217	Human umbilical vein endothelial cells Human aortic endothelial cells	SirT1	(137)




	Human coronary artery endothelial cells		
miR-9	Rats Cardiomyocytes PBMCs	Myocardin	(125-126)
miR-155 and miR-802	CHO cells C2BBel cell line Rat aortic adventitial fibroblasts	AT ₁ R	(106-108)
miR-296-5p, let-7e, hcmv-miR-UL112 and hcmv-miR-UL1	Plasma	IRF-1 MICB	(109)
miR-483-3p	HEK-AT ₁ R HEK-AT ₂ R HASMC HL1-AT ₁ R RASMC RASMC-AT ₁ R RASMC-AT ₂ R Whole heart (C3H NT) Whole heart (C3H TG) Whole heart (C57BL/6 NT) Whole heart (C57BL/6 TG)	AGT ECA-1 ECA2 AT ₂ R	(110)
miR-122	Plasma Mouse Endothelial Cells	rs41318021 polymorphism in the SLC7A1 gene 3'UTR of the L-Arginine Transporter Gene	(113-115)

		<i>SLC7A1</i>	
miR-24	Cardiac endothelial cells Mouse Endothelial cells, cardiomyocytes and cardiac fibroblasts HUVECs and HMVECs Human adrenocortical cell line	GATA2 PAK4 CYP11B1 CYP11B2	(116-118)
miR-124 and miR-135a	Study <i>In silico</i>	Mineralocorticoid receptor <i>NR3C2</i>	(119)
miR-526b, miR-578, miR-34a, miR-34c, miR-765 mir-153-3p and miR-383	Study of Myocardial Infarctions Leiden	Four SNPs located in the arginine vasopressin 1A receptor, bradykinin 2 receptor and thromboxane A2 receptor	(112)
miR-584 and miR-31	Hep3B cells HEK293 cells Genetically hypertensive mice (BPH/2J)	Polymorphism at 11525C in hAGT 3'-UTR	(120, 140, 141)
miR-181a and miR-663	HEK293 cells Plasma	Renin	(104-105)
miR-29b	Mouse heart Cardiac fibroblasts Rat renal medullary epithelial cells	TGF- β 1 Collagen genes	(138-139)
miR-132 and miR-212	Rat heart, aorta and kidney Rat Plasma	AT1R	(111)


	Human artery		
miR-199a-3p	Adipocytes	NLK VAMP3 Cdk7	(143)
miR-326	Plasm	ADIPOR-1 ADIPOR-2 APPL-1	(144)
miR-133a	Plasm	-	(142)

Sprouty-related protein (Spre-1); Vascular Smooth Muscle Cells (VSMCs); Vascular Endothelial Growth Factor (VEGF); Vascular Cell Adhesion Molecule 1 (VCAM-1); Phosphoinositide-3-kinase regulatory subunit 2 (PIK3R2); Phosphatase and tensin homolog (PTEN); Ras Homolog Family Member B (RhoB); Protein sprouty homolog 2 (Sprouty-2); Angiotensin-Converting Enzyme (ACE); Serum Response Factor (SRF); Silent information regulator 1 (SirT1); Angiotensin II Receptor (AT₁R); Transforming Growth Factor (TGF)- β 1; Angiotensinogen (AGT); nemo-like kinase (NLK); vesicle associated membrane protein 3 (VAMP3); cyclin-dependent kinase 7 (Cdk7); adiponectin receptor 1(ADIPOR-1); adiponectin receptor 2 (ADIPOR-2); Adaptor protein, phosphotyrosine interacting with PH domain and leucine zipper 1 (APPL-1)






- Verification of adherence to therapy through daily consultations;
- Referral to multidisciplinary team;
- Conducting ABPM;




- Heart Disease;
- Nephropathy;
- Cerebrovascular Disease

Specialized Investigations




- Sedentary Lifestyle;
- Smoking;
- Alcoholism;
- Obesity;
- Excessive sodium intake;

- Determination 24h urinary sodium ;
- Dietary guidelines;
- Physical exercise;
- Avoid excessive salt and alcohol intake;



- Correct use of medicines;
- Appropriate combinations;
- Adjust dose;



- Primary Aldosteronism;
- Pheochromocytoma;
- Obstructive Sleep Apnea;
- Renal Artery Stenosis;
- Renal Parenchymal Disease;
- Thyroid Disease;

Check Table: Forms of Secondary Hypertension Associated with Resistant Hypertension

